



Commentary

Does homeostasis or disturbance of homeostasis in minimum leaf water potential explain the isohydric versus anisohydric behavior of *Vitis vinifera* L. cultivars?

Jean-Christophe Domec^{1,2,3,4} and Daniel M. Johnson³

¹ University of Bordeaux, Bordeaux Sciences AGRO, UMR 1220 TCEM INRA, 1 Cours du général de Gaulle, 33175 Gradignan Cedex, France; ² Department of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC 27695, USA; ³ Duke University, Durham, NC 27708, USA; ⁴ Corresponding author (jdomec@ncsu.edu)

Received December 31, 2011; accepted January 27, 2012; published online March 15, 2012; handling Editor Danielle Way

Due to the diurnal and seasonal fluctuations in leaf-to-air vapor pressure deficit (D), one of the key regulatory roles played by stomata is to limit transpiration-induced leaf water deficit. Different types of plants are known to vary in the sensitivity of stomatal conductance (g_s) to D with important consequences for their survival and growth. Plants that minimize any increase in transpiration with increasing D have a tight stomatal regulation of a constant minimum leaf water potential (Ψ_{leaf}); these plants are termed as 'isohydric' (Stocker 1956). Plants that have less control of Ψ_{leaf} have been termed as 'anisohydric' (Tardieu and Simonneau 1998). Isohydric plants maintain a constant Ψ_{leaf} by reducing g_s and transpiration under drought stress. Therefore, as drought pushes soil water potential (Ψ_{soil}) below this Ψ_{leaf} set point, the plant can no longer extract water for gas exchange. Anisohydric plants allow Ψ_{leaf} to decrease with rising D , reaching a much lower Ψ_{leaf} in droughted plants relative to well-watered plants (Tardieu and Simonneau 1998), so this strategy produces a gradient between Ψ_{soil} and Ψ_{leaf} that allows gas exchange to continue over a greater decline in Ψ_{soil} . Thus, anisohydric plants sustain longer periods of transpiration and photosynthesis, even under large soil water deficit, and are thought to be more drought tolerant than isohydric species (McDowell 2011).

In practice, the distinctions between isohydric and anisohydric strategies are often not clear (Franks et al. 2007), even among different cultivars of the same species. For example, cultivars of poplar (Hinckley et al. 1994) and grapevine (Schultz 2003, Lovisolo et al. 2010) have been shown to exhibit both contrasting hydraulic behaviors. A third mode of

behavior was also suggested by Franks et al. (2007), in which the difference between soil and midday water potential ($\Psi_{\text{soil}} - \Psi_{\text{leaf}}$) is maintained seasonally constant but Ψ_{leaf} fluctuates in synchrony with soil water availability (isohydrodynamic behavior). The lack of a clear distinction between these two strategies and the complex and variable responses of stomata to D under high and low soil moisture is depicted in two papers in this issue (Rogiers et al. 2012 and Zhang et al. 2012), showing that even typically anisohydric grape (*Vitis vinifera* L.) cultivars (Semillon and Merlot, respectively) may constrain g_s during periods of extremely low Ψ_{soil} . The same individuals can switch from an isohydric-like behavior when transpiration is low to an anisohydric-like behavior with increasing water demand. Interestingly, both studies indicated that classifying species as either isohydric or anisohydric is a simplistic view of stomatal functioning and does not represent well the complex stomatal behavior under drying soil, and Zhang et al. (2012) also reported an isohydrodynamic behavior. Both studies suggested that when soil water is limited, g_s is aimed at protecting the integrity of the hydraulic system, whereas as soil water content increases, stomata regulate transpiration less. The results of Zhang et al. (2012) indicated that under limited soil moisture the decrease in g_s with increasing D was proportional to reference g_s (g_s at $D = 1$ kPa); which is in agreement with the stomata-sensitivity model developed by Oren et al. (1999) for isohydric species (see xeric line in Figure 1A). However, a significant departure from this theoretical model was observed under high soil moisture (see wet and mesic lines in Figure 1B). Similarly, in this issue Rogiers et al. (2012) showed that under




Figure 1. Responses of light-saturated stomatal conductance (g_s) to water vapor pressure deficit (D) as a function of soil moisture (wet, mesic, xeric) in (A) isohydric and (B) anisohydric species as depicted in two papers in this issue (Rogiers et al. 2012 and Zhang et al. 2012). For isohydric species and under saturated light, Oren et al. (1999) showed that the decrease in g_s with increasing D is proportional to g_s at low D . Therefore, the sensitivity (Sens.) of the stomatal response to D can be determined by fitting the data to the functional form: $g_s = g_{s-ref} - Sens. \times \ln(D)$ where g_{s-ref} (designated as reference stomatal conductance) is g_s at $D = 1$ kPa ($\ln D = 0$), and Sens. is the rate of stomatal closure and reflects the sensitivity of g_s to D [$-dg_s/d\ln D$, in $\text{mmol m}^{-2} \text{s}^{-1} \ln(\text{kPa})^{-1}$]. The slope, $-Sens.$ is proportional to g_{s-ref} with the proportionality averaging 0.60 across a wide range of species (Oren et al. 1999). For isohydric species (A), it has been shown that g_{s-ref} and Sens. are both a function of soil moisture and whole-plant hydraulic conductance (K_{plant}) (Domec et al. 2009). For anisohydric species (B), g_{s-ref} and stomatal sensitivity to D can decrease as soil moisture increases and that the same individuals can switch from an anisohydric-like behavior when soil water moisture content is high to an isohydric-like behavior when soil water is low (Rogiers et al. 2012, Zhang et al. (2012). In this issue, Zhang et al. (2012) attributed the departure from an anisohydric to isohydric behavior to whole-plant hydraulic conductance K_{plant} and Rogiers et al. (2012) to the combined effect of K_{plant} and hormonal signaling on g_s by suggesting that the limited soil moisture induced an increase in endogenous ABA, which in turn triggered the increase in stomatal responsiveness to D . To go further, we propose that K_{plant} is driven by change in leaf hydraulic conductance (K_{leaf}) when soil moisture is high, and by root hydraulic conductance (K_{root}) when soil moisture is low and that based on the results from Rogiers et al. (2012), the K_{root} -induced decline in K_{plant} in dry soil acted in concert with ABA production to close stomata.

natural conditions, stomatal sensitivity to D increased under dry soil conditions and that, as in Zhang et al. (2012), stomatal sensitivity to D was proportional to reference g_s . Specifically, Rogiers et al. (2012) demonstrated that reference g_s decreased in most treatments with increasing soil moisture, meaning that

vines exhibited a high stomatal sensitivity to D in very dry soils but not in wet soils (see xeric, mesic and wet lines in Figure 1B). Although Rogiers et al. (2012) also showed that stomatal sensitivity to D increased with reference g_s , their slope was higher than the previously reported generic value of -0.6 based on the Oren et al. (1999) hydraulic model. Rogiers et al. (2012) attributed this departure to the combined effect of leaf hydraulic and hormonal signaling on g_s (Figure 1B). Their results suggested that the limited soil moisture induced an increase in endogenous abscisic acid (ABA), which in turn triggered the increase in stomatal responsiveness to D . The stomatal closure in both studies likely occurred in response to a decrease in leaf or stem hydraulic conductance, but other factors cannot be excluded. Feed-forward behavior of stomata with respect to regulation of Ψ_{leaf} has been attributed to the presence of chemical signals brought to the leaf in the transpiration stream (Davies et al. 1994). Thus, a combination of hydraulic and hormonal signal in some species could be a mechanism allowing some species to switch from an isohydric to anisohydric behavior.

The existence of both isohydric and anisohydric behaviors raises the question of what the costs are for species that operate at more negative Ψ_{leaf} under drought. Compared with isohydric species, anisohydric species maintain carbon assimilation at higher rates as drought intensifies, but this strategy may put anisohydric plants at greater risk of xylem dysfunction if drought is sufficiently intense to push Ψ_{soil} below water potentials that would induce hydraulic failure (Hoffmann et al. 2011, McDowell 2011). Potential drawbacks associated with lower Ψ_{leaf} include a decrease in xylem conductivity due to embolism and a requirement for increased lignin investment in both xylem and leaf structure to resist tension-induced xylem collapse (see Hacke et al. 2001, Blackman et al. 2010). It makes sense that some plants have the potential to switch from an isohydric to anisohydric behavior in order to maximize carbon gain when soil moisture is available and limit plant desiccation when soil moisture is low. Recently, another grape variety, Shiraz, which has been traditionally classified as anisohydric, exhibited isohydric stomatal behavior in dry soils (Collins et al. 2010). These studies raise questions concerning the nature of the mechanisms involved in both stomatal strategies and how the same individuals can switch from one to the other. We can also wonder if such behavior is typical of cultivated grapevines or if it can be extended to lianas in general and perhaps to other plant species.

The underlying mechanism of the so-called 'humidity response', where stomatal conductance decreases with increasing D , remains unknown (Jones 1998, Meinzer 2002). Most of the mechanisms used to explain isohydric behavior are based on stomatal sensitivity to changes in D or soil moisture. However, there are several other potential mechanisms that can help in maintaining seasonal constancy of

plant water status, including the utilization of stored water or the changes in transpiring surface area as shown in Zhang et al. (2012), where increases in whole-plant leaf-specific hydraulic conductance (K_{plant}) were, at least partially, explained by a reduction in the leaf surface during the dry season. Stomata react to prevent embolism at different levels of Ψ_{leaf} , and so the mechanisms controlling stomatal closure also involve differences in hydraulic architecture in the leaves: the onset of leaf vein cavitation makes Ψ_{leaf} drop sharply to maintain the same transpiration rate and thus acts as a signal for stomatal closure (Brodribb and Jordan 2008, Johnson et al. 2009). Evidence that stomata are able to sense and respond rapidly to xylem cavitation, or to changes in K_{plant} associated with dynamics of embolism (Cochard et al. 2002, Domec et al. 2006), implies that embolism in certain plant organs and species, including grapevine, may play a rather different role than was originally believed (Zwieniecki et al. 2000). Chouzouri and Schultz (2005) and more recently, Zufferey et al. (2011) showed that there was a good correlation between drought resistance and vulnerability to embolism in several grapevine cultivars of different geographic origin, and that the decrease in g_s occurred concomitantly with an increase in cavitation events for all varieties, as soil water availability became restricted. Therefore, isohydric and anisohydric behavior under well-watered and water-stressed conditions may be explained by differences in K_{plant} on both a seasonal basis, as soil water varies in time, and on a diurnal basis, as D varies while soil water availability remains nearly constant (Domec et al. 2009). While differences in stomatal behavior in these two strategies may be best explained at the whole-plant level, K_{plant} reflects the conductance of various plant organs (leaves, stems and roots), which can be investigated individually to search for the mechanistic basis of isohydric versus anisohydric behavior (Figure 1A and B). Because of the crucial importance of leaf hydraulic conductance (K_{leaf}) in determining g_s and its sensitivity to water potential (Brodribb and Jordan 2008), variation in leaf hydraulics and its components in response to water availability should receive greater attention in studies aimed at understanding isohydric versus anisohydric behavior (Figure 1A and B). It seems that leaf cavitation plays an important role as a 'hydraulic fuse', thereby limiting leaf transpiration and the propagation of embolism and preserving the integrity of other organs (Chen et al. 2010, Johnson et al. 2011). For example, can we predict that species with a hydraulic design limited by K_{leaf} should be better suited to supply water evenly to the leaf, which should increase stomatal sensitivity to environmental changes (Figure 1A)? Although vine species are characterized by a very efficient water transport system, those species also seem to have developed mechanisms allowing them to refill and repair damaged vessels (Zufferey et al. 2011). As suggested by McDowell (2011), and in this issue by Rogiers

et al. (2012) and Zhang et al. (2012), anisohydric species may be more susceptible than isohydric species to drought-related mortality induced by hydraulic failure. However, it has been reported that in general, anisohydric plants are better adapted to drier environments because of their more cavitation-resistant xylem (McDowell 2011, Hoffmann et al. 2011), or maybe by some (still) unknown refilling mechanisms (Lovisolo et al. 2008). Can we therefore identify xylem structural features that confer morphological advantages, allowing Ψ_{leaf} to approach leaf vein cavitation thresholds without incurring damage from xylem hydraulic failure? To answer such questions, an empirical approach combined with a stomatal hydro-mechanical model is essential (Schultz 2003, Franks et al. 2007). This will provide a better understanding of the complex signals (both chemical and hydraulic) underlying the multi-sensory behavior of stomata.

As suggested by the results of Rogiers et al. (2012), the isohydric and anisohydric stomatal behaviors of grapevine varieties in response to slowly developing water stress imply very different strategies in terms of water use. In countries (mostly in Europe) where irrigation is prohibited when grapes are grown for wine making, varieties that cannot switch from an anisohydric to isohydric behavior may be at risk. Under future climates with potential reductions in precipitation and an increase in the intensity of summer drought, irrigation may play an essential, even if controversial, role in viticulture (Lovisolo et al. 2010), to maintain cultivation of some grapevine varieties for wine production.

Acknowledgments

Danielle Way and two anonymous reviewers are gratefully acknowledged for providing helpful and in-depth comments.

Funding

This commentary was supported by the National Science Foundation (IOS grant 09-20355) and by the Department Of Energy (BER Terrestrial Ecosystem Sciences program, proposal 11-DE-SC-0006700–grant ER65189).

Conflict of interest

None declared.

References

- Blackman, C.J., T.J. Brodribb and G.J. Jordan. 2010. Leaf hydraulic vulnerability is related to conduit dimensions and drought resistance across a diverse range of woody angiosperms. *New Phytol.* 188:1113–1123.
- Brodribb, T.J. and G.J. Jordan. 2008. Internal coordination between hydraulics and stomatal control in leaves. *Plant Cell Environ.* 31:1557–1564.

- Chen, J.-W., Q. Zhang, X.-S. Li and K.-F. Cao. 2010. Gas exchange and hydraulics in seedlings of *Hevea brasiliensis* during water stress and recovery. *Tree Physiol.* 30:876–885.
- Chouzouri, A. and H.R. Schultz. 2005. Hydraulic anatomy, cavitation susceptibility and gas-exchange of several grapevine cultivars of different geographical origin. *Acta Hort.* 689:325–331.
- Cochard, H., L. Coll, X. Le Roux and T. Améglio. 2002. Unraveling the effects of plant hydraulics on stomatal closure during water stress in walnut. *Plant Physiol.* 128:282–290.
- Collins, M.J., S. Fuentes and E.W.R. Barlow. 2010. Partial root-zone drying and deficit irrigation increase stomatal sensitivity to vapour pressure deficit in anisohydric grapevines. *Funct. Plant Biol.* 37:128–138.
- Davies, W.J., F. Tardieu and C.L. Trejo. 1994. How do chemical signals work in plants that grow in drying soil? *Plant Physiol.* 104:309–314.
- Domec, J.-C., F.G. Scholz, S.J. Bucci, F.C. Meinzer, G. Goldstein and R. Villalobos Vega. 2006. Diurnal and seasonal variation in root xylem embolism in neotropical savanna woody species: impact on stomatal control of plant water status. *Plant Cell Environ.* 29:26–35.
- Domec, J.-C., A. Noormets, J.S. King, G. Sun, S.G. McNulty, M.J. Gavazzi, J.L. Boggs and E.A. Treasure. 2009. Decoupling the influence of leaf and root hydraulic conductances on stomatal conductance and its sensitivity to vapor pressure deficit as soil dries in a drained loblolly pine plantation. *Plant Cell Environ.* 32:980–991.
- Franks, P.J., P.L. Drake and R.H. Froend. 2007. Anisohydric but isohydrodynamic: seasonally constant plant water potential gradient explained by a stomatal control mechanism incorporating variable plant hydraulic conductance. *Plant Cell Environ.* 30:19–30.
- Hacke, U.G., J.S. Sperry, W.T. Pockman, S.D. Davis and K.A. McCulloh. 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* 126:457–461.
- Hinckley, T.M., J.R. Brooks, J. Cermák, R. Ceulemans, J. Kucera, F.C. Menzies and D.A. Roberts. 1994. Water flux in a hybrid poplar stand. *Tree Physiol.* 14:1005–1018.
- Hoffmann, W.A., R.M. Marchin, P. Abit and O.L. Lau. 2011. Hydraulic failure and tree dieback are associated with high wood density in a temperate forest under extreme drought. *Glob. Change Biol.* 17:2731–2742.
- Johnson, D.M., K.A. McCulloh, D.R. Woodruff and F.C. Meinzer. 2009. Leaf hydraulic conductance, measured *in situ*, declines and recovers daily: leaf hydraulics, water potential and gas exchange in four temperate and three tropical tree species. *Tree Physiol.* 29:879–887.
- Johnson, D.M., K.A. McCulloh, F.C. Meinzer, D.R. Woodruff and D.M. Eissenstat. 2011. Hydraulic patterns and safety margins, from stem to stomata, in three eastern US tree species. *Tree Physiol.* 31:659–668.
- Jones, H.G. 1998. Stomatal control of photosynthesis and transpiration. *J. Exp. Bot.* 49(Special Issue):387–398.
- Lovisolo, C., I. Perrone, W. Hartung and A. Schubert. 2008. An abscisic acid-related reduced transpiration promotes gradual embolism repair when grapevines are rehydrated after drought. *New Phytol.* 180:642–651.
- Lovisolo, C., I. Perrone, A. Carra, A. Ferrandino, J. Flexas, H. Medrano and A. Schubert. 2010. Drought-induced changes in development and function of grapevine (*Vitis* spp.) organs and in their hydraulic and non-hydraulic interactions at the whole-plant level: a physiological and molecular update. *Funct. Plant Biol.* 37:98–116.
- Meinzer, F.C. 2002. Co-ordination of liquid and vapor phase water transport properties in plants. *Plant Cell Environ* 25:265–274.
- McDowell, N.G. 2011. Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality. *Plant Physiol.* 155:1051–1059.
- Oren, R., J.S. Sperry, G.G. Katul, D.E. Pataki, B.E. Ewers, N. Phillips and K. Schafer. 1999. Survey and synthesis of intra- and interspecific variation in stomatal sensitivity to vapour pressure deficit. *Plant Cell Environ.* 22:1515–1526.
- Rogiers, S.Y., D.H. Greer, J.M. Hatfield, R.J. Hutton, S.J. Clarke, P.A. Hutchinson and A. Somers. 2012. Stomatal response of an anisohydric grapevine cultivar to evaporative demand, available soil moisture and abscisic acid. *Tree Physiol* 32:249–261.
- Schultz, H.R. 2003. Differences in hydraulic architecture account for near-isohydric and anisohydric behaviour of two field-grown *Vitis vinifera* L. cultivars during drought. *Plant Cell Environ.* 26:1393–1405.
- Stocker, O. 1956. Die Abhängigkeit des transpiration von den umwelt-faktoren. In *Encyclopedia of Plant Physiology*. Vol. 3. Ed. W. Ruhland. Springer, Berlin, pp 436–488.
- Tardieu, F. and T. Simonneau. 1998. Variability among species of stomatal control under fluctuating soil water status and evaporative demand: modelling isohydric and anisohydric behaviours. *J. Exp. Bot.* 49:419–432.
- Zhang, Y., R. Oren and S. Kang. 2012. Spatiotemporal variation of crown-scale stomatal conductance in an arid *Vitis vinifera* L. cv. Merlot vineyard: direct effects of hydraulic properties and indirect effects of canopy leaf area. *Tree Physiol* 32:262–279.
- Zufferey, V., H. Cochard, T. Améglio, J.-L. Spring and O. Viret. 2011. Diurnal cycles of embolism formation and repair in petioles of grapevine (*Vitis vinifera* cv. Chasselas). *J. Exp. Bot.* 62:3885–3894.
- Zwieniecki, M.A., L. Hutyra, M.V. Thompson and N.M. Holbrook. 2000. Dynamic changes in petiole specific conductivity in red maple (*Acer rubrum* L.), tulip tree (*Liriodendron tulipifera* L.) and northern fox grape (*Vitis labrusca* L.). *Plant Cell Environ.* 23:407–414.