

## soils & hydrology

# Effects of Timber Harvest on Water Quantity and Quality in Small Watersheds in the Piedmont of North Carolina

Johnny Boggs, Ge Sun, and Steven McNulty

This paired watershed study tested the effects of timber harvest on water quantity and quality in the North Carolina Piedmont physiographic region. Four headwater watersheds at Hill Demonstration Forest (HF1, HF2, HFW1, and HFW2) and two at Umstead Research Farm (UF1 and UF2) were continuously monitored for discharge and water quality from 2007 to 2013. The HF1 and UF1 watersheds were clearcut (treatment), leaving a 15.2-m vegetated riparian buffer around the streams to protect water quality as described in the North Carolina Neuse River Basin Riparian Buffer Rule. HF2 and UF2 were uncut and used as reference watersheds. Merchantable timber was selectively removed from the riparian buffer, reducing tree basal area by 27% in HF1 and 48% in UF1. HF1 and HF2 were nested within HFW1; thus, HFW1 was considered a partial cut where 33% of the watershed area was harvested, and HFW2 was the reference. We found that discharge in treatment watersheds increased dramatically, averaging 240% in HF1 and 200% in UF1 and 40% in HFW1 during the postharvest period, 2011–2013. Total suspended sediment export in the treatment watersheds also increased significantly in HF1 after harvest, probably due to the increase of discharge and movement of in-channel legacy sediment. Stormflow peak nitrate reached its maximum concentration during the first 2 years after harvest in the treatment watersheds and then declined, corresponding to the rapid regrowth of woody and herbaceous plants in the riparian buffer and uplands. We found that 36% of the UF1 streambank trees were blown down but did not cause a measurable increase in mean daily stormflow total suspended sediment concentration. Most buffer tree blowdown occurred during the first few years after a harvest. Bioclassification of benthic macroinvertebrates indicated that stream water quality remained good/fair to excellent in the treatment watersheds after the harvest. We conclude that the temporary increases in discharge were relatively large for the Piedmont region compared with those for other regions in the southeastern United States. However, the increases in channel sediment transport and nutrient exports associated with the hydrologic change did not have a measurable impact on the indicators of aquatic invertebrate community health or bioclassification rankings.

**Keywords:** riparian buffer, best management practices, hydrology, water quality, North Carolina Piedmont

Received September 2, 2014; accepted May 26, 2015; published online August 13, 2015.

**Affiliations:** Johnny Boggs ([jboggs@fs.fed.us](mailto:jboggs@fs.fed.us)), USDA Forest Service, Eastern Forest Environmental Threat Assessment Center, Raleigh, NC. Ge Sun ([gesun@fs.fed.us](mailto:gesun@fs.fed.us)), USDA Forest Service. Steven McNulty ([smcnulty@fs.fed.us](mailto:smcnulty@fs.fed.us)), USDA Forest Service.

**Acknowledgments:** This research was funded by the US Environmental Protection Agency Non-Point Source (NPS) Pollution Control Grant through Section 319(h) of the Clean Water Act, the North Carolina Department of Environment and Natural Resources Division of Water Resources, and the USDA Forest Service Eastern Forest Environmental Threat Assessment Center. We thank the many students and support staff who have been involved with this project over the years for their diligent work in the field and laboratory. We also especially thank William "Bill" Swartley, David Jones, Tom Gerow, and Susan Gale from the North Carolina Forest Service Forestry Nonpoint Source Branch for their project support, guidance, and manuscript review and Will Summer for his field and project development work for this study. Appreciation is expressed to Joe Cox from North Carolina State University and David Schnake from the North Carolina Department of Agriculture and Consumer Services for their partnership and cooperation in providing access to their respective forestlands to conduct this long-term study.

This article uses metric units; the applicable conversion factors are: centimeters (cm): 1 cm = 0.39 in.; meters (m): 1 m = 3.3 ft; square meters (m<sup>2</sup>): 1 m<sup>2</sup> = 10.8 ft<sup>2</sup>; millimeters (mm): 1 mm = 0.039 in.; kilograms (kg): 1 kg = 2.2 lb; milligrams (mg): 1 mg = 0.015 gram; hectares (ha): 1 ha = 2.47 ac.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial reuse, please contact the *Journal of Forestry* at [journal@safnet.org](mailto:journal@safnet.org).

logic and water quality relationship through time between watershed pairs (US Environmental Protection Agency 1993). Preharvest calibration is a major step in assessing and developing predictive models of treatment impacts in a paired watershed design (Swank et al. 2001). The length of the calibration period varies across watersheds due to different controlling factors such as watershed size, soil types, surface cover, and topography (Brooks et al. 2003).

Although the paired watershed approach represents the most rigorous method to quantify forest management effects on water quantity and quality, its applications are often costly and time-consuming. There is also a lack of scientific data and rigorous, long-term watershed-scale forest hydrologic studies in the Piedmont region. In addition, there is still some debate about site-specific design criteria for riparian buffer zones such as type of vegetation management and buffer size. For example, research has shown that when trees in riparian buffers of a certain size are exposed to high winds, blowdown can occur (Grizzell and Wolff 1998). Quantifying the role of evapotranspiration (ET) in watershed response to disturbance in Piedmont watersheds can help with the development of forestry best management practices (BMPs) for reducing stormflow and watershed degradation (Boggs and Sun 2011).

This study seeks to quantify the differences between stream discharge and water quality characteristics (e.g., total suspended sediment, nutrients, and temperature) of forested Piedmont watersheds under undisturbed conditions and clearcuts. We hypothesize that the relationships of water quantity and quality parameters between the treatment watersheds and the reference watersheds will be significantly different from the established relationship of those same watersheds in their uncut state. We also hypothesize that any changes in water quality and quantity will not result in a measurable and sustained change in the benthic bioclassification rankings. Data from this study are useful for addressing land management challenges linked to timber harvesting and other silvicultural activities in the Piedmont region in the southeastern United States.

## Materials and Methods

### Study Sites

We used a standard paired watershed approach in this 6(+) year (2007–2013) monitoring study. Two paired watersheds


were located in the Hill Demonstration Forest (HF) and Umstead Research Farm (UF). The two paired watersheds (treatment watershed HF1 versus reference HF2, treatment watershed UF1 versus reference UF2) and two other watersheds (partial treatment HFW1 versus reference HFW2) are located in the North Carolina Piedmont region (Figure 1). The HFW1 and HFW2 watershed pair was not a part of the original design, but given that the V-notch weirs were already installed in the 1960s as part of another study, we took advantage of the opportunity to monitor them. HF1 and HF2 are nested within HFW1. Therefore, HFW1 is considered a partial treatment watershed where about one-third of the total watershed area was clearcut. HF1, HF2, UF1, and UF2 range from 12 to 29 ha in size with perennial stream channels fitted for stream discharge and water quality monitoring. Dominant overstory species on these sites include red maple (*Acer rubrum*), pignut hickory (*Carya glabra*), mockernut hickory (*Carya tomentosa*), white oak (*Quercus alba*), northern red oak (*Quercus rubra*), American beech (*Fagus grandifolia*), sweetgum (*Liquidambar styraciflua*), tulip poplar (*Liriodendron tulipifera*), sourwood (*Oxydendrum arboreum*), and loblolly pine (*Pinus taeda*). HF1 and HF2 are located in the Flat River Watershed at the North Carolina State University HF in northern Durham County. UF1 and UF2 are located in the Knap of Reeds Watershed at the North Carolina Department of Agriculture and Consumer Services UF in western Granville County. We also monitored hydrology and water quality in the two

larger (HFW1 and HFW2) watersheds at HF. A small portion (~10%) of HFW2 was cut before the beginning of this study, but this did not appear to alter the discharge characteristics or other conditions such that it could not serve as a reference watershed.

HF streams (HF1, HF2, HFW1, and HFW2) are about 1 m wide and 30 cm deep, connected to their narrow floodplain, and have a rocky substrate. These stream channels have steep upland slopes ranging from 15 to 40% with watersheds underlain by soils that have features that are consistent with the Carolina Slate Belt (CSB) region. HF upland soils are defined as well drained with depth to water table of >2 m and tend to function in a similar capacity in the growing season and dormant season. UF streams (UF1 and UF2) are about 2 m wide and 1.5 m deep, are detached from their wide floodplain, and have sandy substrate and gentle upland slopes averaging 7%. UF watersheds are underlain by Triassic Basins (TB) soil characteristics, which are clayey with lower permeability, higher shrink swell characteristics, and thinner soil layers than Carolina Slate Belt soils (US Department of Agriculture 1971). They also generally have a 10-cm thick confining clay layer 30 cm below ground surface, which creates an impermeable condition that results in a perched water table during the dormant season. These features cause variability in how TB soils store, release, and generate water between the growing season and dormant season. TB soils cover about 3.5% of North Carolina land (Cleland et al. 2007) and extend to a small portion of South Carolina.

## Management and Policy Implications

There are three distinct land provinces across North Carolina: the mountains, Piedmont, and coastal plain. Understanding how region-specific watershed hydrology responds to land management and natural disturbances can provide useful information to land managers as they set flow targets needed to maintain ecological integrity in surface waters or to design riparian buffers for water quality protection. We found that after a clearcut timber harvest with a riparian buffer zone around the streams, the percent increases in annual discharge and mean nitrate concentrations tended to be higher in the Piedmont region than in the other two regions. Our study also found that tree blowdown in the riparian buffer was more likely to occur in the Piedmont Triassic region than in the Piedmont Carolina Slate Belt region. In the Triassic region, additional management activities should be part of the preharvest planning process as they may help mitigate windthrow and uprooting of streambanks and improve the overall riparian buffer functions and flow dynamics. For example, land planners or loggers should assess whether soils have Triassic or Slate Belt characteristics to determine if they need to refrain from creating gaps and retain more windfirm species within the riparian buffer. Although there was variation in blowdown occurrences among regions, Triassic soils comprise about 3.5% of North Carolina's total land area, so any statewide implications would be low and recommendations would be localized.



|                      | Hill Demonstration Forest   |     |      |      | Umstead Research Farm  |     |
|----------------------|---|-----|------|------|--|-----|
|                      | HF1   | HF2 | HFW1 | HFW2 | UF1  | UF2 |
| Watershed size (ha)  | 12  | 12  | 29   | 40   | 19   | 29  |
| Stream length (m)    | 300   | 260 | 800  | 960  | 550  | 200 |
| Stand type           | Mixed-pine hardwood   |     |      |      | Mixed-pine hardwood  |     |
| Stand Age (years)    | 35  |     |      |      | 70   |     |
| Slope (%)            | 13  |     |      |      | 7  |     |
| Geologic Features    | Carolina Slate Belt   |     |      |      | Triassic Basin   |     |
| Dominate Soil Series | Tatum and Appling   |     |      |      | Helena   |     |
| Soil characteristics | Non expansive clays, no perched water, deep soils, discharges water slowly throughout the year due to large amounts of stored water in bedrock and topographic control. |     |      |      | Expansive clays, perched water, thin soils, discharges water slowly in growing season when soils are dry with an in-active confining clay layer, and fast in dormant season when soils are wet with an active confining layer. |     |

**Figure 1.** Aerial views of paired study watersheds and descriptive attributes.

Additional details on stream channels can be found in Boggs et al. (2013) and Dreps et al. (2014).

### Riparian Vegetation Surveys

To characterize vegetation composition along a 10% reach of the stream study area, 4 152-m<sup>2</sup> vegetation survey plots in HF1, 6 plots in HF2, 10 plots in UF1, and 4 plots in UF2 were established. Stem count, dbh of overstory trees, and canopy cover in each plot were measured annually following protocols outlined in the Carolina Vegetation Survey (Peet et al. 1998). Percent canopy cover was measured at plot center with hemispherical photography every year during the growing season. Six 1-m<sup>2</sup> subplots were also established in each plot to estimate percent groundcover. Visual observations for riparian buffer sediment breakthroughs or overland flow were assessed at least monthly or after large ( $>25$  mm) storm events along the entire reach and width of

the channel. Breakthroughs were determined to be one of the following: overland and sediment flowing through the riparian buffer to the channel; only overland flow moving through the riparian buffer to the channel; or evidence of overland flow or sediment moving into the riparian buffer but being dispersed before reaching the stream (Rivenbark and Jackson 2004). Field inspections in April 2013 and August 2013 revealed considerable blowdown in one of the treatment watersheds (UF1). Thus, additional vegetation surveys were taken to determine the number and diameter size of standing and windthrown stream edge trees. Stream edge trees were defined as trees having roots exposed in the stream channel. All blowdown trees had tip-up mounds with some mounds being as large as 3 m in diameter (Figure 2). If a stream edge tree had a broken top, it was not counted as blowdown.

### Harvest

The entire watershed areas at both study sites were clearcut harvested using typical rubber tire-mounted logging equipment. Logging on UF1 took place between July 7 to Sept. 8, 2010, and logging on HF1 occurred from Nov. 29, 2010 to Jan. 19, 2011. In each clearcut harvest, a 15.2-m riparian buffer was retained on each side of the stream. High-value trees (trees of  $\geq 35.6$  cm dbh for pine and  $\geq 40.6$  cm for hardwood) were harvested from the riparian buffer as allowed by the Neuse River Basin Riparian Buffer Rule (NRR). Hand felling of high-value and merchantable timber within inner zone 1 (0–3 m from the stream bank) and outer zone 1 (3–9.1 m from the stream bank) of the riparian buffer was done as outlined in the NRR. On each study site, additional BMPs were deployed to prevent sedimentation and water quality pollution and to comply with the North Carolina Forest



**Figure 2.** Tip-up mound of blowdown of stream edge tree in UF1 at Umstead Research Farm, Granville County, North Carolina. Photo courtesy of Johnny Boggs, USDA Forest Service, August 2013. Pictured in photo: Neil Williams, Forestry Technician, USDA Forest Service.

Practices Guidelines Related to Water Quality. For example, trees were skidded to the log deck without crossing the stream channel, and slash was redistributed across the upland and skid trails to limit soil disturbance. Site preparation for replanting included one aerial herbicide application, a bladed fireline around the watershed boundary, and a low-intensity site preparation burn that was initiated from outside of the riparian buffer. A fireline was not put around the riparian buffer. The site preparation process and hand planting operation occurred between June 2011 and January 2012 in HF1 and between July 2011 and January 2012 in UF1. Loblolly pine was planted in HF1, and shortleaf pine (*Pinus echinata*) was planted in UF1.

### Stream Discharge and Water Quality Measurements

A 2-H flume was used as the flow control structure at the outlet of HF1, HF2, UF1, and UF2, and a 90° V-notch weir was used at the outlet of HFW1 and HFW2. A Sigma 900 Max water sampler with a depth sensor was used to measure and log discharge data every 10 minutes in units of liters/second. This unit was then converted to mm to normalize the watershed discharge data and to make the discharge data comparable to precipitation. Discharge data in this

article are reported in mm (see Supplemental Figure S1<sup>5</sup> to convert mm back to liters). Precipitation was measured in an open area with a HOBO Datalogging Rain Gauge RG3 at HF and UF. Water quality concentrations and exports were quantified from grab and storm-based samples. Grab water samples were collected at least biweekly under baseflow conditions. The Sigma sampler was programmed to trigger based on an increase in the flow rate of change (e.g., 1.1 liters/second). Storm-based samples were collected on a stratified sampling program, intensive sampling during rising limb (6 samples in 1 hour) and less intense sampling during recession limb (6 samples over 6–10 hours) of the hydrograph. To avoid the potential to overemphasize one limb of the hydrograph (or to interpolate between measured times), a time-weighted mean concentration for each constituent was computed and then flow weighted concentrations were determined.

Water quality parameters included total suspended sediment (TSS), total organic carbon (TOC), ammonium-nitrogen ( $\text{NH}_4\text{-N}$ ), nitrate-nitrogen ( $\text{NO}_3\text{-N}$ ), total phosphorus (TP), total Kjeldahl nitrogen (TKN), stream temperature, and a macroinvertebrate bioclassification or biotic index. Samples were preserved with sulfuric

acid to pH of <2. Water samples collected from the field were kept at 3.6° C before analysis. Constituents from each water sample were determined at the North Carolina State University Soil Science Analytical Laboratory using standard methods (Greenburg 1992). The changes in water quality parameters were initially measured in mg liter<sup>-1</sup> during both stormflow and baseflow conditions. These values were then multiplied by discharge volume to determine outputs expressed as kg ha<sup>-1</sup> month<sup>-1</sup> and kg ha<sup>-1</sup> year<sup>-1</sup>. Total organic nitrogen (TON) equals TKN minus  $\text{NH}_4\text{-N}$ . Total nitrogen (TN) equals TKN plus  $\text{NO}_3\text{-N}$ . Stream temperature data were logged every 10 minutes using HOBO Pro v2 water temperature sensors.

During the preharvest period, two benthic surveys were completed in all six watersheds but only covered the nongrowing season because of limited sampling time. Seven postharvest surveys were taken in all watersheds and covered both growing and non-growing seasons. Surveys were taken across seasons to capture differences in the life cycle of benthic macroinvertebrates, seasonal discharge, and climate variability. Benthic macroinvertebrate surveys were completed following the semiquantitative methods outlined by the North Carolina Department of Environmental and Natural Resources (2012) Division of Water Resources, Biological Assessment Unit Qual4 method, for which a kick net, sweep net, leaf pack, and visual samples were collected from each stream. Benthic macroinvertebrate samples were field sorted and sent to Watershed Science, LLC, to be identified to the lowest possible taxonomic class. To rate water quality condition, a bioclassification class (excellent, good, good/fair, fair, or poor) for small streams (i.e., <4.6 m wide) was assigned to each survey, which was based on the average values from Ephemeroptera, Plecoptera, and Trichoptera (EPT) taxa richness and biotic index (Lenat 1993, 2014). EPT taxa richness is the number of Ephemeroptera, Plecoptera, and Trichoptera taxa in a sample. Generally, the higher the taxa richness is, the better the water quality condition. The EPT taxon captures the full range of water quality conditions through time, whereas baseflow and stormflow samples do not fully capture conditions between samples. A mixture of both water chemistry and benthic surveys

<sup>5</sup> Supplementary data are available with this article at <http://dx.doi.org/10.5849/jof.14-102>.

probably offers the best data to assess water quality condition.

The dominant trophic category or functional feeding group (FFG) percentages (i.e., collector gather, scraper collector, obligate scraper, shredder, predator, collector filters, and omnivore scavenger) for each macroinvertebrate species was calculated by assigning a semiquantitative value to the occurrence class: rare = 1, common = 3, and abundant = 10. The FFG percentages were used in conjunction with bioclassification and other indicators to provide additional information about the treatment effect and to assess process-level attributes about the aquatic ecosystem (Rawer-Jost et al. 2004).

## Data Processing and Analysis

The experimental design consisted of a pair of watersheds (reference and treatment), a calibration or preharvest period, a treatment (i.e., tree harvest), and a postharvest period (Swank et al. 2001). In the preharvest phase (2007–2010), discharge, TSS, and the water quality parameters from the paired watersheds were calibrated.

To calibrate the watersheds, a set of linear relationship/models ( $y = mx + b$ ) between daily discharge and monthly TSS and nutrient concentrations and exports from each pair were generated with all  $P < 0.05$ . Postharvest reference watershed data were then put into that linear model to predict what the treatment watershed trend would be if the harvest had not occurred. Postharvest modeled treatment data and postharvest measured reference data were compared to evaluate the treatment effect. Our paired watersheds did not have identical soil and vegetative composition and stormflow discharge. Thus, calibrating the reference watershed to the treatment watershed provided a more accurate assessment of treatment effects on discharge, water quality data, and cause-effect relationships compared with referencing the treatment watershed directly. Our 32-month calibration period developed a robust enough model for these small catchments to account for any changes or variation in climate and other factors during the postharvest period (US Environmental Protection Agency 1993). Wilm (1944, 1948) developed the following equation to determine the minimum length of calibration time required for a watershed to predict treatment effects with a reasonable level of certainty (in our study, measured

and modeled discharge was within 2 SD of each other)

$$k = \frac{s_{y,x}^2 F}{d^2}$$

where  $k$  equals the number of observations from each of the two data sets,  $s_{y,x}^2$  is the SE of the estimate ( $y$ ) in mm/month, the  $F$  statistic equals  $F[2 + [F/(k - 1)]]$ , and  $d$  equals the smallest noteworthy change in monthly runoff. Based on Wilm's equation calibration, watersheds (HF1 and HF2) could be calibrated in 10 months at  $\alpha = 0.05$  and 32 months at  $\alpha = 0.01$  (Boggs et al. 2008).

We analyzed mean annual ( $n = 3$ ) TSS and nutrient concentrations and exports across preharvest and postharvest periods using a  $t$ -test (SAS Institute, Inc. 2011). The  $t$ -test was selected, and the significance level was set to  $\alpha \leq 0.05$  to determine which group values (measured versus modeled) were statistically different from each other. Significant differences statements are  $\alpha \leq 0.05$ . Storm parameters were derived from a constant slope separation method where water is discharged from a watershed in excess of 1.1 mm/day as described by Hewlett and Hibbert (1967). The constant slope value was applied to the separation analysis during 13 to 44 storms when at least 15–20 mm of measured rainfall occurred. Slope separation was terminated when the total volume of discharge exceeded the baseflow discharge. The average separation analysis lasted 21.2 (SE, 3.3 hours) hours during the nongrowing season (November to April) and 12.9 (2.2) hours during the growing season (May to October).

## Results

### Vegetation Changes in the Riparian Buffer

The riparian buffer vegetation for the preharvest period was consistent with a basal area range found in the closed-canopy Piedmont mixed pine-hardwood riparian forests (Supplemental Table S1; Figure 1). According to the NRR, trees can be removed from the riparian buffer during logging. Selective removal of trees from the riparian buffer reduced HF1 pine and hardwood overstory basal areas by 24 and 28%, respectively. Tree removal from the riparian buffer reduced UF1 pine and hardwood overstory basal areas by 50 and 46%, respectively. The reference watershed total overstory basal area increased 13% from 2009 to 2013 in

both HF2 (30.5–34.5  $m^2 ha^{-1}$ ) and UF2 (36.9–41.8  $m^2 ha^{-1}$ ).

There were two events of stream edge tree blowdown in UF1. One was documented on Apr. 17, 2013, in which 11% of stream edge canopy trees blew down, and the other on Aug. 28, 2013, when an additional 25% of stream edge canopy trees blew down. Trees in UF1 that were above the mean riparian buffer dbh blew down more often than trees below the mean buffer dbh, with hickory and oak species experiencing 100% blowdown within the buffer (Table 1). Pine and sweetgum experienced 50% or greater blowdown when trees were above the mean dbh. When there was more than one tree present, tulip poplar was the most windfirm tree species (i.e., those least likely to be blown down during high wind events). Tree blowdown was also documented on Mar. 14, 2014, in treatment watershed (HF1) at the HF site. This documented blowdown was outside of the water quality monitoring period for this study, and none of the wind-thrown trees were located at the stream edge.

Percent groundcover shifted in treatment watersheds from leaf litter dominated in 2009 (preharvest, Figure S2a) to a mixture of woody, herbaceous, and leaf litter dominated from 2011 to 2013 (postharvest, Figure S2b). Percent groundcover in reference watersheds (HF2 and UF2) remained dominated by leaf litter from 2009 to 2013 (Figures S2c and d). UF2 herbaceous groundcover increased in 2013 compared with that in other years (Figure S2d) probably due to an opening in the canopy structure around one of the riparian buffer survey plots. The higher SE (7.8%) of herbaceous groundcover in the UF site than in other plots is reflective of localized canopy openness. We did not observe sediment flowing overland and through the riparian buffer to the channel during the monitoring period at the UF or HF site.

### Precipitation and Discharge


Cumulative precipitation amounts for the preharvest and postharvest periods were 3,818 and 3,256 mm at HF and 3,460 mm and 3,266 mm at UF, respectively. Measured and modeled cumulative discharges were similar during the preharvest period but increased dramatically during the postharvest period (Tables 2 and 3; Figure 3 and Supplemental Figure S3). Measured postharvest discharge increased above modeled postharvest discharge by 263, 264, and 192% in HF1, by 45, 45, and 37% in HFW1, and by 248,

**Table 1. Percentage of stream edge trees that blew down in treatment watershed, UF1.**

| Common name    | Species (scientific name)      | Stream edge tree dbh range (cm) | Percent of stream edge trees that blew down, larger or equal to the mean dbh (%) | Percent of stream edge trees that blew down, smaller than mean dbh (%) | Total no. of all trees tallied on stream edge, both standing and blownover | No. of stream edge trees that blew down, larger or equal to the mean dbh | No. of stream edge trees that blew down, smaller than mean dbh | Total no. of stream edge trees that DID NOT blowover |
|----------------|--------------------------------|---------------------------------|--|--|--|--|--|--|
| American beech | <i>Fagus grandifolia</i>       | 18–18                           | — <sup>a</sup>   | 0  | 2  | —  | 0  | 2  |
| Blackgum       | <i>Nyssa sylvatica</i>         | 30–30                           | —  | 0  | 2  | —  | 0  | 2  |
| Elm            | <i>Ulmus</i> spp.              | 20                              | —  | 0  | 1  | —  | 0  | 1  |
| Hickory        | <i>Carya</i> spp.              | 20–33                           | 100  | 0  | 4  | 1  | 0  | 3  |
| Ironwood       | <i>Carpinus caroliniana</i>    | 18                              | —  | 0  | 1  | —  | 0  | 1  |
| Oak            | <i>Quercus</i> spp.            | 18–76                           | 100  | 0  | 6  | 4  | 0  | 2  |
| Pine           | <i>Pinus</i> spp.              | 38–53                           | 50   | —  | 4  | 2  | —  | 2  |
| Red cedar      | <i>Juniperus virginiana</i>    | 20                              | —  | 100  | 1  | —  | 1  | 0  |
| Red maple      | <i>Acer rubrum</i>             | 13–20                           | —  | 25   | 4  | —  | 1  | 3  |
| Sourwood       | <i>Oxydendrum arboreum</i>     | 10–15                           | —  | 0  | 7  | —  | 0  | 7  |
| Sweetgum       | <i>Liquidambar styraciflua</i> | 13–76                           | 65   | 38   | 25   | 11   | 3  | 11   |
| Sycamore       | <i>Platanus occidentalis</i>   | 33                              | 0  | —  | 1  | 0  | —  | 1  |
| Tulip poplar   | <i>Liriodendron tulipifera</i> | 15–79                           | 20   | 0  | 8  | 1  | 0  | 7  |

Blowdown percentage is separated into above and below mean diameter at breast height (dbh) to test whether larger trees were more vulnerable to blowdown than smaller trees. Mean stream edge tree dbh was 33 cm.

<sup>a</sup>—, tree size not present at stream edge in the riparian buffer.



**Figure 3. Daily precipitation, cumulative measured discharge with harvest, and modeled discharge without harvest in treatment watersheds over monitoring period (a) HF1, (b) HFW1, (c) UF1. Rectangle indicates tree harvest period; HF1 and HFW1 were harvested Nov. 29, 2010 to Jan. 19, 2011 and UF1 was harvested July 7, 2010 to Sept. 8, 2010.**

**Table 2. Measured and modeled discharge, precipitation, percent ET, and percent discharge of precipitation during preharvest and postharvest period in a timber harvest study at Hill Demonstration Forest, Durham County, NC.**

| Year                                      | n<br>(days) | Period             | HF1 measured | HF1 modeled | HF2 measured | HFW1 measured | HFW1 modeled | HFW2 measured | Precipitation HF |
|---|-------------|--------------------|--------------|-------------|--------------|---------------|--------------|---------------|------------------|
| (mm)                                      |             |                    |              |             |              |               |              |               |                  |
| 2007                                      | 105         | Preharvest         | 7            | 13          | 17           | 20            | 29           | 8             | 152              |
| 2008                                      | 366         | Preharvest         | 117          | 113         | 142          | 151           | 199          | 171           | 1,197            |
| 2009                                      | 365         | Preharvest         | 209          | 190         | 236          | 262           | 256          | 257           | 1,358            |
| 2010                                      | 365         | Preharvest         | 170          | 190         | 236          | 286           | 236          | 226           | 1,096            |
| 2011                                      | 18          | Preharvest         | 6            | 4           | 5            | 7             | 5            | 1             | 14               |
|   |             | Total              | 508          | 510         | 637          | 725           | 725          | 663           | 3,817            |
|   |             | ET (%) preharvest  | 87           | 87          | 83           | 81            | 81           | 83            |                  |
| 2011                                      | 347         | Postharvest        | 247          | 68          | 88           | 175           | 121          | 62            | 1,002            |
| 2012                                      | 366         | Postharvest        | 251          | 69          | 89           | 180           | 124          | 60            | 1,057            |
| 2013                                      | 365         | Postharvest        | 327          | 112         | 142          | 291           | 212          | 191           | 1,198            |
|   |             | Total              | 826          | 249         | 318          | 646           | 457          | 313           | 3,256            |
|   |             | ET (%) postharvest | 75           | 92          | 90           | 80            | 86           | 90            |                  |
| <i>Percent discharge of precipitation</i> |             |                    |              |             |              |               |              |               |                  |
| 2007                                      | 105         | Preharvest         | 4            | 9           | 11           | 13            | 19           | 5             |                  |
| 2008                                      | 366         | Preharvest         | 10           | 9           | 12           | 13            | 17           | 14            |                  |
| 2009                                      | 365         | Preharvest         | 15           | 14          | 17           | 19            | 19           | 19            |                  |
| 2010                                      | 365         | Preharvest         | 16           | 17          | 22           | 26            | 22           | 21            |                  |
| 2011                                      | 18          | Preharvest         | 41           | 27          | 35           | 48            | 36           | 10            |                  |
|   |             | Mean               | 17           | 15          | 19           | 24            | 22           | 14            |                  |
| 2011                                      | 347         | Postharvest        | 25           | 7           | 9            | 17            | 12           | 6             |                  |
| 2012                                      | 366         | Postharvest        | 24           | 7           | 8            | 17            | 12           | 6             |                  |
| 2013                                      | 365         | Postharvest        | 27           | 9           | 12           | 24            | 18           | 16            |                  |
|   |             | Mean               | 25           | 8           | 10           | 20            | 14           | 9             |                  |

Percent ET is estimated as (total precipitation – total discharge)/total precipitation × 100 over the preharvest and postharvest monitoring periods. Percent discharge of precipitation equals (precipitation/discharge × 100). HF1 and HFW1 harvest ended on Jan. 19, 2011. Measured discharge is cumulative flow values from the pressure transducer. Modeled discharge is cumulative flow values from the linear model that was developed during the calibration period to determine what discharge would be if the clear cut had not occurred. HF1 and HFW1, treatment watersheds; HF2 and HFW2, reference watersheds.

**Table 3. Measured and modeled discharge, precipitation, percent ET, and percent discharge of precipitation during preharvest and postharvest period in a timber harvest study at Umstead Research Farm, Granville County, NC.**

| Year                                      | n<br>(days) | Period             | UF1 measured<br>discharge | UF1 modeled<br>discharge | UF2 measured<br>discharge | Precipitation<br>UF |
|---|-------------|--------------------|---------------------------|--------------------------|---------------------------|---------------------|
| (mm)                                      |             |                    |                           |                          |                           |                     |
| 2007                                      | 105         | Preharvest         | 2                         | 1                        | 3                         | 198                 |
| 2008                                      | 366         | Preharvest         | 128                       | 122                      | 164                       | 1,249               |
| 2009                                      | 365         | Preharvest         | 281                       | 286                      | 372                       | 1,307               |
| 2010                                      | 251         | Preharvest         | 125                       | 145                      | 189                       | 705                 |
|   |             | Total              | 537                       | 554                      | 728                       | 3,460               |
|   |             | ET (%) preharvest  | 84                        | 84                       | 79                        |                     |
| 2010                                      | 114         | Postharvest        | 52                        | 15                       | 23                        | 223                 |
| 2011                                      | 365         | Postharvest        | 181                       | 52                       | 78                        | 985                 |
| 2012                                      | 366         | Postharvest        | 261                       | 82                       | 117                       | 967                 |
| 2013                                      | 365         | Postharvest        | 376                       | 155                      | 208                       | 1,091               |
|   |             | Total              | 870                       | 304                      | 427                       | 3,266               |
|   |             | ET (%) postharvest | 75                        | 91                       | 88                        |                     |
| <i>Percent discharge of precipitation</i> |             |                    |                           |                          |                           |                     |
| 2007                                      | 105         | Preharvest         | 1                         | 0.3                      | 1                         |                     |
| 2008                                      | 366         | Preharvest         | 10                        | 10                       | 13                        |                     |
| 2009                                      | 365         | Preharvest         | 22                        | 22                       | 28                        |                     |
| 2010                                      | 251         | Preharvest         | 18                        | 21                       | 27                        |                     |
|   |             | Mean               | 13                        | 13                       | 17                        |                     |
| 2010                                      | 114         | Postharvest        | 23                        | 7                        | 10                        |                     |
| 2011                                      | 365         | Postharvest        | 18                        | 5                        | 8                         |                     |
| 2012                                      | 366         | Postharvest        | 27                        | 8                        | 12                        |                     |
| 2013                                      | 365         | Postharvest        | 34                        | 14                       | 19                        |                     |
|   |             | Mean               | 26                        | 9                        | 12                        |                     |

Percent ET is estimated as (total precipitation – total discharge)/total precipitation × 100 over the preharvest and postharvest monitoring periods. Percent discharge of precipitation equals (precipitation/discharge – 100). UF1 harvest ended on Sept. 8, 2010. Measured discharge is cumulative flow values from the pressure transducer. Modeled discharge is cumulative flow values from the linear model that was developed during the calibration period to determine what discharge would be if the clearcut had not occurred. UF1, treatment watershed; UF2, reference watershed.

**Table 4. Mean values for stormflow hydrologic characteristics of treatment watersheds with harvest (measured) and without harvest (modeled) during the postharvest period.**

| Watershed                    | Season     | No. of storms | Event duration            | Beginflow                | Peak rate                 | Time peak   | Total discharge          | Baseflow                 | Stormflow                | Precipitation             | D/P ratio                |
|------------------------------|------------|---------------|---------------------------|--------------------------|---------------------------|-------------|--------------------------|--------------------------|--------------------------|---------------------------|--------------------------|
|                              |            |               | hr                        | ..... (mm/day).....      |                           | hr          | .(mm/storm) .            |                          |                          |                           |                          |
| HF1 measured with harvest    | Growing    | 40            | 11.00 (1.32) <sup>a</sup> | 0.70 (0.13) <sup>a</sup> | 9.43 (3.44)               | 2.75 (0.58) | 1.65 (0.53) <sup>a</sup> | 0.53 (0.11) <sup>a</sup> | 1.12 (0.44)              | 24.90 (1.98)              | 0.05 (0.01) <sup>a</sup> |
| HF1 modeled without harvest  | Growing    | 40            | 6.37 (0.82) <sup>a</sup>  | 0.19 (0.04) <sup>a</sup> | 5.11 (0.90)               | 2.02 (0.41) | 0.61 (0.25) <sup>a</sup> | 0.13 (0.04) <sup>a</sup> | 0.49 (0.21)              | 22.50 (1.74)              | 0.02 (0.01) <sup>a</sup> |
| HFW1 measured with harvest   | Growing    | 44            | 10.38 (1.08) <sup>a</sup> | 0.56 (0.09)              | 7.82 (2.18) <sup>a</sup>  | 2.48 (0.36) | 1.36 (0.39)              | 0.43 (0.09)              | 0.92 (0.31)              | 23.71 (1.67) <sup>a</sup> | 0.05 (0.01)              |
| HFW1 modeled without harvest | Growing    | 33            | 7.49 (0.97) <sup>a</sup>  | 0.48 (0.06)              | 2.59 (0.59) <sup>a</sup>  | 1.86 (0.15) | 0.59 (0.18)              | 0.27 (0.06)              | 0.33 (0.12)              | 18.77 (1.72) <sup>a</sup> | 0.03 (0.01)              |
| UF1 measured with harvest    | Growing    | 36            | 17.21 (1.68) <sup>a</sup> | 0.35 (0.07) <sup>a</sup> | 41.60 (7.94) <sup>a</sup> | 3.52 (0.51) | 7.17 (1.48) <sup>a</sup> | 0.69 (0.11) <sup>a</sup> | 6.48 (1.39) <sup>a</sup> | 25.86 (2.76)              | 0.24 (0.04) <sup>a</sup> |
| UF1 modeled without harvest  | Growing    | 33            | 9.96 (1.72) <sup>a</sup>  | 0.07 (0.02) <sup>a</sup> | 14.12 (3.97) <sup>a</sup> | 2.47 (0.52) | 2.69 (0.87) <sup>a</sup> | 0.23 (0.06) <sup>a</sup> | 2.47 (0.82) <sup>a</sup> | 21.08 (2.54)              | 0.09 (0.02) <sup>a</sup> |
| HF1 measured with harvest    | Nongrowing | 23            | 23.48 (8.05)              | 0.79 (0.18) <sup>a</sup> | 5.24 (1.60)               | 5.64 (1.14) | 2.11 (0.68) <sup>a</sup> | 0.86 (0.19) <sup>a</sup> | 1.25 (0.51)              | 23.05 (2.51)              | 0.07 (0.01) <sup>a</sup> |
| HF1 modeled without harvest  | Nongrowing | 23            | 10.75 (1.30)              | 0.32 (0.05) <sup>a</sup> | 3.65 (0.59)               | 4.31 (0.88) | 0.72 (0.13) <sup>a</sup> | 0.28 (0.07) <sup>a</sup> | 0.44 (0.08)              | 20.33 (1.74)              | 0.03 (0.00) <sup>a</sup> |
| HFW1 measured with harvest   | Nongrowing | 19            | 14.77 (2.19)              | 0.64 (0.05)              | 3.74 (0.66)               | 5.32 (1.12) | 1.50 (0.46)              | 0.70 (0.18)              | 0.80 (0.28)              | 20.22 (2.32)              | 0.06 (0.01)              |
| HFW1 modeled without harvest | Nongrowing | 13            | 14.68 (2.55)              | 0.55 (0.05)              | 3.03 (0.77)               | 5.76 (1.21) | 1.34 (0.44)              | 0.66 (0.20)              | 0.68 (0.25)              | 15.43 (2.94)              | 0.08 (0.02)              |
| UF1 measured with harvest    | Nongrowing | 20            | 25.47 (2.34)              | 0.48 (0.11) <sup>a</sup> | 33.01 (6.62) <sup>a</sup> | 6.29 (1.07) | 9.03 (1.92) <sup>a</sup> | 1.33 (0.29)              | 7.70 (1.72) <sup>a</sup> | 22.97 (2.75)              | 0.33 (0.04) <sup>a</sup> |
| UF1 modeled without harvest  | Nongrowing | 19            | 21.38 (2.46)              | 0.24 (0.07) <sup>a</sup> | 13.58 (2.79) <sup>a</sup> | 6.13 (1.08) | 4.65 (0.99) <sup>a</sup> | 0.79 (0.16)              | 3.85 (0.85) <sup>a</sup> | 21.78 (2.77)              | 0.19 (0.03) <sup>a</sup> |

Measured stormflow parameters are values from the pressure transducer. Modeled stormflow parameters are values from the linear model that was developed during the calibration period to determine what stormflow would be if the clearcut had not occurred. Standard errors are in parentheses.

<sup>a</sup>t-test to determine significant difference ( $P < 0.05$ ) between measured with harvest versus modeled without harvest within watershed and season.

218, and 143% in UF1 from 2011, 2012, and 2013, respectively. Although precipitation totals were similar between preharvest and postharvest periods, the percent discharge of precipitation decreased from preharvest to postharvest period in the reference watersheds due to fewer (8 versus 3) intense storms ( $\geq 25.4$  mm/hour) and smaller discharge events. These factors probably led to greater amounts of deep seepage and water storage during postharvest than during preharvest in the reference watersheds.

Stormflow characteristics changed after harvest in all treatment watersheds (Table 4). Measured growing season stormflow duration with harvest was significantly higher than that of modeled growing season stormflow duration without harvest in all watersheds. Measured growing and nongrowing season beginflow and discharge to precipitation ratios in the treatment watershed were significantly higher than modeled beginflow and discharge to precipitation ratio in both HF1 and UF1. This indicated that growing and nongrowing season soils were wetter after harvest in HF1 and UF1. Measured peak rate of stormflow was significantly higher than the modeled peak rate in UF1 during both growing and nongrowing seasons and

both were significantly higher in HFW1 only during the growing season. Measured total discharge increased significantly above the modeled total discharge in HF1 and UF1 during both the growing and nongrowing seasons. In UF1, the growing and nongrowing season stormflow increased significantly.

#### Mean Annual, Peak Nitrate, and Daily Stormflow Water Quality Concentration

Mean annual measured preharvest TSS and nutrient concentrations were not significantly different from mean annual modeled preharvest concentrations in either watershed (Table 5). All values were within background levels for forests or near the detection limits. Mean annual measured TP, TN, and TON concentrations were highest in HF1 compared with those for HFW1 and UF1. UF1 had a higher mean annual measured TOC concentration than the other treatment watersheds.

Mean annual measured NO<sub>3</sub>-N concentration during the postharvest period was statistically higher than mean annual modeled NO<sub>3</sub>-N concentration in both HF1 and UF1 with NO<sub>3</sub>-N reaching 0.13 mg liter<sup>-1</sup> in HF1 and 0.45 mg liter<sup>-1</sup> in UF1

(Table 5). Mean annual measured TP concentration was significantly higher than the modeled value in UF1 during the postharvest period. Mean annual measured TSS, TOC, NO<sub>3</sub>-N, TN, and TON concentrations were higher in UF1 than in HF1 and HFW1.

The stream nitrate concentration peaked during the postharvest period at 4.9 mg liter<sup>-1</sup> in HF1, at 4.5 mg liter<sup>-1</sup> in HFW1, and at 6.9 mg liter<sup>-1</sup> in UF1 (Supplemental Figure S4). By the mid-2012 postharvest period, NO<sub>3</sub>-N peak concentrations in the treatment watersheds were near preharvest peak concentrations. The mean daily stormflow TSS concentrations in the treatment watersheds were similar between preharvest, postharvest, and blowdown periods (Supplemental Figure S5).

#### Mean Annual Export of TSS and Nutrients

Mean annual measured TSS and nutrient export values during the preharvest period were not significantly different from mean annual modeled preharvest exports in all watersheds (Table 6). In the postharvest period, all mean annual measured constituents were higher than the mean annual modeled constituents in all treatment water-

**Table 5. Mean annual measured and modeled TSS and nutrient concentrations during preharvest and postharvest periods in the treatment watersheds.**

| Period                              | Constituents       | HF1                      |                           | HFW1        |             | UF1                       |                           |
|-------------------------------------|--------------------|--------------------------|---------------------------|-------------|-------------|---------------------------|---------------------------|
|                                     |                    | Measured                 | Modeled                   | Measured    | Modeled     | Measured                  | Modeled                   |
| .....(mg liter <sup>-1</sup> )..... |                    |                          |                           |             |             |                           |                           |
| Preharvest                          | TSS                | 36.8 (0.02)              | 35 (0.04)                 | 27.9 (9.7)  | 26.7 (5.2)  | 35.7 (9.3)                | 33.8 (6.4)                |
|                                     | TOC                | 6.1 (6.4)                | 6.2 (2.9)                 | 5.7 (0.9)   | 5.7 (0.5)   | 10.0 (0.96)               | 9.5 (0.96)                |
|                                     | NH <sub>4</sub> -N | 0.03 (0.3)               | 0.04 (0.4)                | 0.01 (0.02) | 0.01 (0.03) | 0.01 (0.007)              | 0.01 (0.005)              |
|                                     | NO <sub>3</sub> -N | 0.01 (0.01)              | 0 (0.01)                  | 0.02 (0.06) |             | 0.03 (0.013)              | 0.01 (0.002)              |
|                                     | TP                 | 0.08 (0.005)             | 0.07 (0.00)               | 0.06 (0.02) | 0.06 (0.01) | 0.07 (0.009)              | 0.06 (0.01)               |
|                                     | TN                 | 0.71 (0.004)             | 0.69 (0.002)              | 0.52 (0.13) | 0.50 (0.08) | 0.66 (0.05)               | 0.67 (0.05)               |
|                                     | TON                | 0.67 (0.06)              | 0.64 (0.004)              | 0.5 (0.12)  | 0.48 (0.06) | 0.62 (0.049)              | 0.59 (0.04)               |
| Postharvest                         | TSS                | 31.1 (0.06)              | 34.3 (0.01)               | 23.8 (0.7)  | 23.2 (1.2)  | 33.6 (0.8)                | 28.6 (2.3)                |
|                                     | TOC                | 4.6 (0.3)                | 5.5 (1.1)                 | 4.7 (0.8)   | 5.2 (0.5)   | 15.1 (3.9)                | 8.2 (0.3)                 |
|                                     | NH <sub>4</sub> -N | 0.08 (0.4)               | 0.03 (0.4)                | 0.02 (0.03) | 0.01 (0.01) | 0.04 (0.005)              | 0.06 (0.036)              |
|                                     | NO <sub>3</sub> -N | 0.13 (0.01) <sup>a</sup> | 0.00 (0.002) <sup>a</sup> | 0.06 (0.05) |             | 0.45 (0.156) <sup>a</sup> | 0.01 (0.005) <sup>a</sup> |
|                                     | TP                 | 0.09 (0.04)              | 0.07 (0.00)               | 0.07 (0.02) | 0.06 (0.01) | 0.08 (0.001) <sup>a</sup> | 0.06 (0.004) <sup>a</sup> |
|                                     | TN                 | 0.81 (0.01)              | 0.64 (0.004)              | 0.52 (0.08) | 0.44 (0.08) | 1.37 (0.20)               | 0.82 (0.17)               |
|                                     | TON                | 0.6 (0.07)               | 0.6 (0.02)                | 0.43 (0.06) | 0.43 (0.07) | 0.89 (0.17)               | 0.60 (0.03)               |

The predictive model was not good enough to develop a reliable modeled value (i.e., probability statistic = 0.43 for NO<sub>3</sub>-N in HFW1 and 0.62 for NH<sub>4</sub>-N in UF1). Measured concentrations are mean values from water samples collected during baseflow and stormflow. Modeled concentrations are mean values from the linear model that was developed during the calibration period to determine what load would be if the clearcut had not occurred. HF1 and HFW1 were harvested from Nov. 29, 2010, to Jan. 19, 2011, and UF1 was harvested July 7, 2010, to Sept. 8, 2010. Standard errors are in parentheses.

<sup>a</sup>t-test to determine significance ( $P < 0.05$ ) between measured and modeled values within the watershed and period.

**Table 6. Mean annual measured and modeled TSS and nutrient loads during preharvest and postharvest periods in the treatment watersheds.**

| Period  | Constituents       | HF1                      |                          | HFW1                     |                          | UF1                      |                          |
|---|--------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
|   |                    | Measured                 | Modeled                  | Measured                 | Modeled                  | Measured                 | Modeled                  |
| .....(kg ha <sup>-1</sup> yr <sup>-1</sup> )..... |                    |                          |                          |                          |                          |                          |                          |
| Preharvest  | TSS                | 74.2 (23.9)              | 73.4 (18.5)              | 82.5 (29.1)              | 82.0 (15.6)              | 93.4 (27.4)              | 85.2 (23.5)              |
|   | TOC                | 9.4 (1.3)                | 9.3 (1.2)                | 13.9 (1.9)               | 13.8 (1.4)               | 22.2 (7.3)               | 22.2 (7.9)               |
|   | NH <sub>4</sub> -N | 0.02 (0.01)              | 0.02 (0.01)              | 0.013 (0.01)             | 0.013 (0.01)             | 0.013 (0.003)            |                          |
|   | NO <sub>3</sub> -N | 0.003 (0.001)            | 0.004 (0.001)            | 0.03 (0.01)              |                          | 0.03 (0.02)              | 0.03 (0.001)             |
|   | TP                 | 0.16 (0.06)              | 0.15 (0.04)              | 0.17 (0.04)              | 0.17 (0.03)              | 0.19 (0.09)              | 0.19 (0.08)              |
|   | TN                 | 1.19 (0.39)              | 1.17 (0.23)              | 1.37 (0.35)              | 1.36 (0.33)              | 1.56 (0.66)              | 1.56 (0.56)              |
|   | TON                | 1.17 (0.39)              | 1.14 (0.23)              | 1.33 (0.35)              | 1.32 (0.32)              | 1.52 (0.67)              | 1.52 (0.56)              |
| Postharvest                                       | TSS                | 94.4 (14.8) <sup>a</sup> | 31.4 (9.6) <sup>a</sup>  | 59.8 (13.9)              | 44.5 (3.3)               | 84.5 (25.9)              | 36.9 (7.0)               |
|   | TOC                | 13.8 (2.2) <sup>a</sup>  | 4.3 (0.6) <sup>a</sup>   | 11.3 (1.8)               | 7.8 (0.8)                | 32.7 (4.1) <sup>a</sup>  | 9.9 (1.8) <sup>a</sup>   |
|   | NH <sub>4</sub> -N | 0.27 (0.12)              | 0.01 (0.002)             | 0.07 (0.03)              | 0.006 (0.001)            | 0.09 (0.02)              |                          |
|   | NO <sub>3</sub> -N | 0.68 (0.33)              | 0.001 (0.001)            | 0.19 (0.1)               |                          | 1.14 (0.46)              | 0.04 (0.02)              |
|   | TP                 | 0.31 (0.1)               | 0.06 (0.01)              | 0.21 (0.1)               | 0.11 (0.02)              | 0.22 (0.06)              | 0.09 (0.02)              |
|   | TN                 | 2.62 (0.26) <sup>a</sup> | 0.72 (0.14) <sup>a</sup> | 1.28 (0.18) <sup>a</sup> | 0.71 (0.07) <sup>a</sup> | 3.13 (0.56) <sup>a</sup> | 0.89 (0.24) <sup>a</sup> |
|   | TON                | 1.67 (0.21) <sup>a</sup> | 0.71 (0.14) <sup>a</sup> | 1.03 (0.17)              | 0.67 (0.07)              | 1.89 (0.40) <sup>a</sup> | 0.79 (0.15) <sup>a</sup> |

The predictive model was not good enough to develop reliable modeled value (i.e., probability statistic = 0.43 for NO<sub>3</sub>-N in HFW1 and 0.62 for NH<sub>4</sub>-N in UF1). Measured loads are mean values from water samples collected during baseflow and stormflow multiplied by discharge. Modeled loads are mean values from linear model that was developed during the calibration period to determine what load would be if the clearcut had not occurred. HF1 and HFW1 were harvested Nov. 29, 2010 to Jan. 19, 2011, and UF1 was harvested July 7, 2010 to Sept. 8, 2010. Standard errors are in parentheses.

<sup>a</sup>t-test to determine significance ( $P < 0.05$ ) between measured and modeled values within the watershed and period.

sheds. However, mean annual measured TN was the only water quality parameter that was significantly higher than the mean annual modeled value across all treatment watersheds. The measured TSS load increased significantly from the modeled TSS in HF1. Although not significant, measured TSS load also increased from modeled TSS in HFW1 and UF1. Mean annual measured TOC and TON increased significantly above mean annual modeled values in both HF1 and UF1. Modeled postharvest loads were lower than modeled preharvest loads for all constituents in all treatment watersheds because percent

discharge of precipitation was smaller during the postharvest period than during the preharvest period.

#### Benthic Macroinvertebrates

Preharvest water quality bioclassification ranged from good to excellent in all treatment watersheds (Table 7). Postharvest bioclassification in harvested sites ranged from good/fair to excellent, whereas those in the reference sites ranged from fair to excellent. In the preharvest period, mean non-growing season EPT taxa richness and biotic index values were not significantly different between any treatment and reference water-

shed pair. In the postharvest period, both mean growing season and nongrowing season EPT taxa richness increased (improved) significantly in HFW1 compared with that for HFW2 and in UF1 compared with that for UF2. The biotic index decreased or improved significantly in HFW1 compared with that in HFW2 during the nongrowing season. The dominant FFG or trophic category was similar between treatment and reference watersheds during the postharvest period (Figure 4). For example, the highest FFG percentage was for shredders in the HF1 and HF2 pair (29 and 27% of the total macroinvertebrates in the samples, respec-

**Table 7. EPT taxa richness, biotic index, and bioclassification in treatment and reference watersheds during preharvest and postharvest periods.**

| Watershed        | Index             | Preharvest   |            |                 |            |           | Postharvest   |           |               |           |              |                         |                         |
|------------------|-------------------|--------------|------------|-----------------|------------|-----------|---------------|-----------|---------------|-----------|--------------|-------------------------|-------------------------|
|                  |                   | January 2010 | April 2010 | Mean nongrowing | March 2011 | July 2011 | February 2012 | July 2012 | February 2013 | June 2013 | January 2014 | Mean growing            | Mean nongrowing         |
| HF1 (treatment)  | EPT taxa richness | 13           | 16         | 14.5 (1.5)      | 9          | 8         | 11            | 9         | 16            | 7         | 12           | 8.0 (0.6)               | 12.0 (1.5)              |
| HF2 (reference)  | EPT taxa richness | 21           | 17         | 19.0 (2.0)      | 14         | 10        | 14            | 12        | 12            | 9         | 15           | 10.3 (0.9)              | 13.8 (0.6)              |
| HFW1 (treatment) | EPT taxa richness | 24           | 22         | 23.0 (1.0)      | 20         | 11        | 20            | 12        | 20            | 11        | 14           | 11.3 (0.3) <sup>a</sup> | 18.5 (1.5) <sup>a</sup> |
| HFW2 (reference) | EPT taxa richness | 20           | 17         | 18.5 (1.5)      | 17         | 2         | 11            | 4         | 13            | 7         | 11           | 4.3 (1.5) <sup>a</sup>  | 13.0 (1.4) <sup>a</sup> |
| UF1 (treatment)  | EPT taxa richness | 16           | 12         | 14.0 (2.0)      | 19         | 10        | 20            | 11        | 14            | 10        | 12           | 10.3 (0.3) <sup>a</sup> | 16.3 (1.9) <sup>a</sup> |
| UF2 (reference)  | EPT taxa richness | 18           | 10         | 14.0 (4.0)      | 10         | 0         | 12            | 1         | 12            | 4         | 10           | 1.7 (1.2) <sup>a</sup>  | 11.0 (0.6) <sup>a</sup> |
| HF1 (treatment)  | Biotic index      | 4.5          | 3.3        | 3.9 (0.60)      | 5.2        | 4.0       | 4.8           | 4.2       | 4.2           | 3.6       | 4.1          | 3.9 (0.19)              | 4.6 (0.26)              |
| HF2 (reference)  | Biotic index      | 3.8          | 3.0        | 3.4 (0.40)      | 4.9        | 3.2       | 4.8           | 3.7       | 4.7           | 3.8       | 3.9          | 3.6 (0.18)              | 4.6 (0.22)              |
| HFW1 (treatment) | Biotic index      | 3.9          | 3.4        | 3.7 (0.25)      | 4.1        | 3.3       | 4.0           | 3.8       | 3.7           | 3.6       | 3.9          | 3.6 (0.15)              | 3.9 (0.08) <sup>a</sup> |
| HFW2 (reference) | Biotic index      | 4.1          | 2.8        | 3.5 (0.65)      | 4.6        | 6.3       | 4.2           | 4.5       | 4.4           | 3.0       | 4.3          | 4.6 (0.95)              | 4.4 (0.08) <sup>a</sup> |
| UF1 (treatment)  | Biotic index      | 4.8          | 4.8        | 4.8 (0.00)      | 5.4        | 4.8       | 4.7           | 4.9       | 4.1           | 4.1       | 5.1          | 4.6 (0.27)              | 4.8 (0.28)              |
| UF2 (reference)  | Biotic index      | 4.5          | 4.0        | 4.3 (0.25)      | 5.4        | 6.6       | 4.5           | 5.4       | 6.3           | 4.6       | 5.2          | 5.5 (0.59)              | 5.3 (0.36)              |
| HF1 (treatment)  | Bioclassification | Good         | Excellent  |                 | Good/fair  | Excellent | Good          | Excellent | Excellent     | Excellent | Excellent    |                         |                         |
| HF2 (reference)  | Bioclassification | Excellent    | Excellent  |                 | Good       | Excellent | Good          | Excellent | Good          | Excellent | Excellent    |                         |                         |
| HFW1 (treatment) | Bioclassification | Excellent    | Excellent  |                 | Excellent  | Excellent | Excellent     | Excellent | Excellent     | Excellent | Excellent    |                         |                         |
| HFW2 (reference) | Bioclassification | Excellent    | Excellent  |                 | Good       | Fair      | Excellent     | Good      | Good          | Excellent | Good         |                         |                         |
| UF1 (treatment)  | Bioclassification | Good         | Good       |                 | Good/fair  | Good      | Good          | Good      | Excellent     | Excellent | Good         |                         |                         |
| UF2 (reference)  | Bioclassification | Good         | Excellent  |                 | Good/fair  | Fair      | Good          | Good/fair | Fair          | Good      | Good         |                         |                         |

Criteria for North Carolina biotic index for small streams (<4 m wide): excellent, <4.3; good, 4.3–5.1; good/fair, 5.2–5.8; fair, 5.9–6.9; poor, >6.9. In the postharvest period, the mean nongrowing season includes March 2011, February 2012, February 2013, and January 2014 and the growing season included July 2011, July 2012, and June 2013. Standard errors are in parentheses.

<sup>a</sup>t-test to determine significance ( $P < 0.05$ ) between mean treatment versus mean reference EPT (Ephemeroptera [mayfly], Plecoptera [stonefly], and Trichoptera [caddisfly]) taxa richness and biotic index values of each watershed pair (HF1 versus HF2, HFW1 versus HFW2, and UF1 versus UF2) during preharvest and postharvest periods and across the nongrowing season or growing season.

tively) and the HFW1 and HFW2 pair (25 and 26%, respectively). Collector gatherers FFG was the highest in the UF1 and UF2 pair (21 and 33%, respectively).

### Stream Temperature

Preharvest monthly maximum stream temperature values were 24.8° C in HF1, 22.4° C in HF2, 25.8° C in HFW1, 24.5° C in HFW2, 23.5° C in UF1, and 24.1° C in UF2. After the harvest, monthly maximum stream temperature spiked during the first two growing seasons in HFW1 and during the first growing season only in UF1 (Supplemental Figure S6a and b). Unexpectedly, a spike in temperature was not observed in the other treatment watershed, HF1, during the postharvest period.

### Discussion


#### Discharge Response to Harvesting

Understanding regional site-specific water resource responses to timber removal and other land disturbances can help land managers set flow targets to reduce stormflow and watershed degradation and manage general streamflow dynamics. When ET is reduced through clearcut harvesting, discharge has been shown to increase by 370 mm (60%) in the mountains (Johnson and Kovner 1954), 91 mm (99%) in the coastal plain of North Carolina (Amatya et al. 2006), and 250 mm (55%) in the Piedmont

of South Carolina (Williams et al. 2000) during the 1st year. During the 1st year after harvest in our study, we found that discharge increased more dramatically after forest removal compared with the values in these studies. Discharge increased 180 mm (263%) over modeled discharge rates in HF1 and 129 mm (248%) over modeled discharge rates in UF1. We believe this large increase in annual discharge postharvest is driven primarily by the type of topography and climatic conditions found in our study catchments and their effects on ET. ET was estimated to be about 80% of the annual water budget for our study sites. In contrast to the results for the Piedmont, Sun et al. (2002) reported that annual ET was 70% of precipitation for a coastal plain pine plantation and 47% of precipitation in a hardwoods watershed at Coweeta in the mountains of North Carolina. Post and Jones (2001) reported that ET in mountainous regions in the United States ranged from about 36 to 50% of the water budget. Thus, when our watersheds were clearcut, a large response in discharge was expected because the tree hydrologic ET pump was removed from the system.

The effect of the disturbance on discharge was less when a smaller portion of our Piedmont watershed was disturbed. For example, HFW1 experienced 33% tree removal from the watershed (HF1 experi-

enced 92% timber removal) that resulted in an increase in discharge of 54 mm (45%) during the first year after harvest. On a 10% forest removal basis ( $54 \text{ mm} \times 10/33\% = 16.4 \text{ mm year}^{-1}$  in HFW1 and  $180 \text{ mm year}^{-1} \times 10/92\% = 19.6 \text{ mm}$  in HF1), the uncut portion of HFW1 dampened the effect on discharge from the cut area, HF1, during year 1 by 20% (i.e.,  $19.6 \text{ mm} - 16.4 \text{ mm}/16.4 \text{ mm}$ ). However, the capacity to buffer against the effects of a clearcut on streamflow may depend on the distance of the disturbance from the stream channel. Based on a simulation study using Visualizing Ecosystems for Land Management Assessments (VELMA), Abdelnour et al. (2011) reported that a 20% clearcut area near the catchment divide resulted in an average annual discharge increase of 53 mm, whereas a 20% clearcut near the stream resulted in an average annual discharge increase of 92 mm. Stednick (1996) reported that when 20% or less of a watershed was harvested, it was unlikely that any increase in annual discharge would be detected. In addition, an increase in peak flow may be undetectable in harvested areas of less than 29% (Grant et al. 2008) or 32% of removal of the basal area (Bent 1994). We detected an increase in peak rate in all treatment watersheds with the largest differences between measured versus modeled occurring in the growing season.



**Figure 4. Mean functional feeding group percentages in treatment versus reference (HF1 versus HF2, HFW1 versus HFW2, and UF1 versus UF2) watersheds during the postharvest period, 2011–2013.**

### Water Quality Response to Harvesting

The significant increases in  $\text{NO}_3\text{-N}$  concentrations in both HF1 and UF1 after harvest were below levels considered harmful to biological integrity (Binkley et al. 1999, Boggs et al. 2013) but were generally higher than values found in the North Carolina mountains (Clinton 2011), Alabama coastal plain (Lockaby et al. 1994), or South Carolina coastal plain (Askew and Williams 1986) or Piedmont (Williams et al. 2000). Quantifying nutrient values in the North Carolina Piedmont after a harvest may not only help to document stream quality responses to disturbances but can also provide further context for managing water resources through improved planning and treatment strategies across regions. Nitrate peaks in our study were probably driven by a reduction in plant nitrogen uptake that left the highly mobile nitrate anions vulnerable to flushing during storm events (Hornberger

et al. 1994). UF1 had considerably more nitrate spikes than HF1 and HFW1 (Supplemental Figure S4) because of the rapid lateral and increased stormflow that occur in these TB soils. UF1 stormflow, peakflow, and total discharge increased significantly in the postharvest period in both growing and nongrowing seasons, increasing the potential to quickly mobilize accumulated nitrate to the stream channel. Increases in peak  $\text{NO}_3\text{-N}$  were delayed by 6–8 months from the time of harvest. This delay is not uncommon as time is needed for ammonium to accumulate in the soil after reduced demand by the trees, and time is also needed for nitrifying populations to increase with the increasing soil ammonium. In addition, time is needed for root systems and slash to decay and release additional N back into the soil. However, despite the pulse increases,  $\text{NO}_3\text{-N}$  levels remained low. By mid-2012, the regrowth of woody and herbaceous

plants is what probably reduced stormflow peak  $\text{NO}_3\text{-N}$  concentrations to preharvest levels in all treatment watersheds. HF1 demonstrated a greater capacity to retain nitrate compared with that for HF1 and UF1 because a much smaller percentage of HF1 was disturbed. Although the UF1 site exhibited the highest peak  $\text{NO}_3\text{-N}$  concentrations over a relatively short period, the TB soils found on the UF1 site only comprise 3.5% of North Carolina's total land area (Cleland et al. 2007), so any impacts would be localized.

Some of the measured water quality loads were not significantly different from modeled loads during the postharvest period because discharge and climate variability increased with increasing exports (Table 6). Despite this statistical limitation, it is still informative to discuss the annual load patterns because they indicate the amount of nutrient impact on the ecosystem. Mean annual measured TN load was the only parameter that increased significantly above mean annual modeled load values in all treatment watersheds. Mean annual measured TP, TOC, and TSS loads increased 1- to 2-fold above mean annual modeled loads across treatments. These increases are consistent with effects of timber removal on sediment and nutrient yields from headwater catchments (Arthur et al. 1998, Fraser et al. 2012). In addition, changes in TP export tend to follow changes in TSS export given that phosphorus often attaches to small particles of sediment (Brady 1990). The increases in TSS and nutrient exports in the treatment watersheds were probably driven by the large increase in discharge in response to the clearcut harvest (Swank and Vose 1994, McBroom et al. 2008) and movement of in-channel legacy sediment during pre-harvest (Boggs et al. 2013) and postharvest periods (J.L. Boggs, USDA Forest Service, unpubl. data, Sept. 8, 2010 to Dec. 31, 2013). Given that postharvest discharge and  $\text{NO}_3\text{-N}$  concentration spikes declined each year and corresponded to increased ET and plant growth, TSS, ammonium, and other nutrient exports will probably match pre-harvest levels within the next 5 years. Overall, our results showing that temporary increases in discharge were accompanied by increased in-channel sediment transport and nutrient exports but did not measurably impact indicators of aquatic invertebrate communities align with those of other timber harvesting studies (Table 8).

The macroinvertebrate bioclassification

**Table 8. Comparison of postharvest discharge, sediment, and nitrate in paired watershed timber harvesting studies.**

| Study                             | Treatment                                    | Location                                | Discharge<br>(mm yr <sup>-1</sup> )   | Sediment<br>.. (mg liter <sup>-1</sup> ) | NO <sub>3</sub> -N<br>.. | Sediment<br>.. (kg ha <sup>-1</sup> yr <sup>-1</sup> ) | NO <sub>3</sub> -N<br>.. | Conclusions  |
|-----------------------------------|--|---|---------------------------------------|--|--------------------------|--|--------------------------|--|
| This study                        |  |   |                                       |  |                          |  |                          |  |
| HF2                               | Control                                      | North Carolina/<br>Piedmont             | 105                                   | 29                                       | 0.003                    | 41   | 0.002                    | Temporary increases in discharge were accompanied by increased in-channel sediment transport and nutrient exports but were not sufficiently disruptive to impact aquatic life and ecological integrity.    |
| HFW2                              | Control                                      |   | 105                                   | 22                                       | 0.001                    | 33   | 0.02                     |  |
| UF2                               | Control                                      |   | 130                                   | 34                                       | 0.29                     | 56   | 0.36                     |  |
| HF1, 15.2-m buffer                | Clearcut with buffer                         |   | 280                                   | 31                                       | 0.13                     | 94   | 0.67                     |  |
| HFW1, 15.2-m buffer               | Clearcut with buffer                         |   | 215                                   | 24                                       | 0.06                     | 61   | 0.19                     |  |
| UF1, 15.2-m buffer                | Clearcut with buffer                         |   | 262                                   | 33                                       | 0.48                     | 101  | 1.37                     |  |
| Fraser et al. (2012)              | Control                                      | Georgia/Piedmont                        |                                       |  |                          | 418  |                          |  |
| 12–21-m buffer                    | Clearcut with buffer                         |   |                                       |  |                          | 1,067  |                          | SMZ greatly reduced the water quality effects of clearcutting.   |
| Clinton 2011                      | Control                                      | North Carolina/<br>Mountains            |                                       | 16                                       | 0.04                     |  |                          |  |
| 10-m buffer                       | Clearcut with buffer                         |   | 32                                    | 0.05                                     |                          |  |                          |  |
| 30-m buffer                       | Clearcut with buffer                         |   |                                       | 8  | 0.02                     |  |                          |  |
| No buffer                         | Clearcut without<br>buffer                   |   |                                       | 24                                       | 0.12                     |  |                          |  |
| McBroom et al. (2008)             | ...SW3, SW5, SW8                             | Texas/East                              | 14                                    |  |                          | 42   |                          |  |
|                                   | ...SW2, SW4, SW9                             | Clearcut/conventional<br>prep with BMPs | 78                                    |  |                          | 111  |                          |  |
|                                   | ...SW1, SW6, SW7                             | Clearcut/intensive<br>prep with BMPs    | 98                                    |  |                          | 225  |                          |  |
| Williams et al. (2000)            | Kenamore One                                 | South Carolina/<br>Piedmont             | 458                                   | 114                                      | 0.03                     | 522  | 0.16                     | Increases in discharge were accompanied by increases in sediment and nutrient exports. Sediment loads without BMPs can be 10-fold higher than loads with BMPs according to Hewlett (1979).                 |
| Holley Springs,<br>12.2-m buffer  | Clearcut with BMPs                           |   | 709                                   | 98                                       | 0.08                     | 894  | 0.58                     |  |
| Ramsey Bridge,<br>12.2-m buffer   | Clearcut/herbicide<br>with BMPs              |   | 677                                   | 95                                       | 0.09                     | 647  | 0.60                     |  |
| Kenamore Two,<br>12.2-m buffer    | Clearcut/mechanical<br>and thin with<br>BMPs |   | 902                                   | 90                                       | 0.14                     | 802  | 0.81                     |  |
| Arthur et al. (1998) <sup>a</sup> | Watershed A                                  | Kentucky/Cumberland<br>Plateau          | 490                                   |  | 1.0                      | 30   | 0.24                     | Harvested watersheds had higher discharge and concentrations of nitrate and other nutrients compared with those for the control. Regrowth of vegetation reduced nutrient losses within 3 years of harvest. |
|                                   | Watershed B, 15.2-m<br>buffer                | Clearcut with BMPs                      | 600                                   |  | 1.8                      | 250  | 1.91                     |  |
|                                   | Watershed C                                  | Clearcut without<br>BMPs                | 650                                   |  | 2.0                      | 300  | 2.18                     |  |
| Kochenderfer et al.<br>(1997)     | Haddix                                       | Clearcut with BMPs                      | West<br>Virginia/Allegheny<br>Plateau |  | 1.45                     | 58   |                          | Sediment loads returned to preharvest levels by the 3rd postharvest year.  |

<sup>a</sup> Sediment and nutrient values were averaged across all postharvest years.

rankings indicated that stream quality remained good/fair to excellent in the treatment watersheds during the postharvest period (Table 7). However, some reference watersheds had biclassification scores that ranked from fair to a maximum of good/fair. These reference watershed scores were often linked to low discharge periods during the growing season that drove down the presence of certain indicator aquatic species. For example, the UF2 July 2011 survey had the lowest EPT taxa richness and was the lowest discharge month (0.13 mm) compared with any other month. The UF2 biclassification for July 2011 was fair, whereas the UF1 biclassification was good and had a measured

discharge of 2.3 mm. UF1 without harvest discharge (based on a model from the calibration period) would have been 0.11 mm. This suggests that the additional discharge in UF1 after the harvest may have improved stream habitat for benthic macroinvertebrates beyond what would have been observed without harvest. However, the dominant FFG (i.e., collector gatherers in UF and shredders in HF) in the treatment and reference watersheds remained the same, and the distribution of FFG in the watershed pairs was similar after harvest (Figure 4). Jackson et al. (2007) also found that macroinvertebrate groups did not decline significantly after harvest. This suggests that

any increase or fluctuations in discharge or nutrient concentrations probably did not pose any risk to water quality, despite differences in bioclassification scores between our watershed pairs during certain surveys.

### Riparian Buffer Vegetation

Our study is consistent with findings from previous work indicating that blowdown occurs during the first few years after a harvest under normal weather events or patterns (Moore 1977, Jackson et al. 2007). In UF1, 36% of stream edge trees blew down during two windstorm events in the third postharvest year. There was no measurable change in UF1 mean daily TSS concentra-

tions in stormflow water samples compared with those in treatment watersheds without blowdown, HF1 and HFW1 (Supplemental Figure S5). Grizzel and Wolff (1998) found a similar result where 33% of buffer trees were windthrown but did not produce a significant amount of additional sediment to the headwater stream channel. They also noted that any sediment delivered to streams from windthrown trees was small relative to the amount stored in the channel. There appears to be some degree of riparian buffer resiliency to the windthrow disturbance that protected water quality in UF1. The resiliency is possibly the result of the increase in herbaceous and woody vegetation growth as HF1 herbaceous plant coverage increased from 0.8% preharvest to 21% postharvest, and UF1 herbaceous plant coverage increased from 2.1 to 28% over the same time period (Figure S2). This additional coverage and plant root structure could provide groundcover protection and soil stability in the riparian buffer (Wynn and Mostaghimi 2006). Streambank failure could occur in the future in UF1 because the edge function of the riparian buffer had been compromised, which may cause excessive sedimentation to occur. Therefore, forestry BMPs and measures should always be implemented to prevent windthrow and uprooting of streambanks to ensure that the riparian buffer function is not compromised.

Different tree blowdown patterns between UF1 and HF1 were probably caused by differences in tree size, species, and soil characteristics at our study sites (Steinblums et al. 1984, Sinton et al. 2000, Steil et al. 2009). The UF1 riparian buffer trees were larger (30 cm) than the HF1 riparian buffer trees (23.8 cm). Trees in UF1 that were above the mean riparian buffer dbh blew down more often than trees below the mean dbh (Table 1), suggesting that larger trees were less windfirm than smaller trees. For trees above a mean riparian buffer dbh of 33 cm and in cases where more than one tree was present, the order for most windfirm species to least windfirm species appears to be tulip poplar > sweetgum > pine spp. > hickory spp. = oak spp. Although there was variation in blowdown occurrences among regions (Triassic versus Carolina Slate Belt), TB soils comprise about 3.5% of North Carolina's total land area, so any statewide implications would be low and recommendations would be site-specific.

## Conclusions

Our 6-year study has resulted in important findings in forest hydrology, nutrient exports, and vegetated riparian buffer functions in Piedmont watersheds. Among similar studies in the region, this paired watershed study provides a complete assessment of hydrology and water quality responses to timber harvesting. We conclude that forest vegetation removal plays a more significant role in affecting water balances and mean and peak nitrate concentrations in this region than in the mountains and coastal plains. However, overall stream water quality in the Piedmont was not negatively affected by increases in discharge, nutrient loading, and stream edge tree blowdown. The knowledge gained from this project will be useful to land managers. It should provide a better understanding of how Piedmont watersheds store, release, and discharge water and nutrients across growing and dormant seasons, how riparian buffers function, and how to apply the most appropriate timber harvest management practices for protecting water resources across regions.

## Literature Cited

- ABDELNOUR, A., M. STIEGLITZ, F. PAN, AND R. MCKANE. 2011. Catchment hydrological responses to forest harvest amount and spatial pattern. *Water Resour. Res.* 47(9):W09521.
- AMATYA, D.M., R.W. SKAGGS, C.D. BLANTON, AND J.W. GILLIAM. 2006. Hydrologic and water quality effects of harvesting and regeneration of a drained pine forest. P. 538–550 in *Hydrology and management of forested wetlands, Proc. of the international conference*. ASABE Publ. 701P0406, American Society of Agricultural and Biological Engineers, St. Joseph, MI.
- ARTHUR, M.A., G.B. COLTHRAP, AND D.L. BROWN. 1998. Effects of best management practices on forest stream water quality in eastern Kentucky. *J. Am. Water Resour. Assoc.* 34: 412–417.
- ASKEW, G.R., AND T.M. WILLIAMS. 1986. Water quality changes due to site conversion in coastal South Carolina. *South. J. Appl. For.* 10: 134–136.
- AUST, W.M., M.B. CARROLL, M.C. BOLDING, AND C.A. DOLLOFF. 2011. Operational forest stream crossings effects on water quality in the Virginia Piedmont. *J. Appl. For.* 35(3):123–130.
- BENT, G.C. 1994. Effects of timber cutting on runoff to Quabbin Reservoir, central Massachusetts. P. 187–196 in *Proc., effects of human-induced changes on hydrologic systems*, Marston, R.A., and V.R. Hasfurther (eds.). American Water Resources Association, Bethesda, MD.
- BINKLEY, D., H. BURNHAM, AND H.L. ALLEN. 1999. Water quality impacts of forest fertilization with nitrogen and phosphorus. *For. Ecol. Manage.* 121(3):191–213.
- BOOGS, J.L., G. SUN, W. SUMMER, S.G. McNULTY, W. SWARTLEY, AND E. TREASURE. 2008. Effectiveness of streamside management zones on water quality: Pretreatment measurements. P. 1–6 in *AWRA summer specialty conference; 2008 June 30–July 2*. USDA For. Serv., Southern Research Station, Asheville, NC.
- BOOGS, J.L., AND G. SUN. 2011. Urbanization alters watershed hydrology in the Piedmont of North Carolina. *Ecohydrology* 4(2):256–264.
- BOOGS, J.L., G. SUN, D.G. JONES, AND S.G. McNULTY. 2013. Effect of soils on water quantity and quality in Piedmont forested headwater watersheds of North Carolina. *J. Am. Water Resour. Assoc.* 49(1):132–150.
- BRADY, N. 1990. *The nature and properties of soils*, 10th ed. Macmillan Publishing, New York. 550 p.
- BROOKS, K.N., P.F. FFOLLIOTT, H.M. GREGERSEN, AND L.F. DEBANO. 2003. *Hydrology and the management of watersheds*, 3rd ed. Iowa State University Press, Ames, IA. 574 p.
- BROWN, T.C., AND D. BINKLEY. 1994. *Effect of management on water quality in North American forests*. USDA For. Serv., Gen. Tech. Rep. RM-248, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 27 p.
- BROWN, T.C., M.T. HOBBINS, AND J.A. RAMIREZ. 2008. Spatial distribution of water supply in the conterminous United States. *J. Am. Water Resour. Assoc.* 44:1474–1487.
- CLELAND, D.T., J.A. FREEOUF, J.E. KEYS JR., G.J. NOWACKI, C. CARPENTER, AND W.H. McNAB. 2007. *Ecological subregions: Sections and subsections of the conterminous United States [1: 3,500,000]* [CD-ROM], Sloan, A.M. (cartog.). USDA For. Serv., Gen. Tech. Rep. WO-76D, Washington, DC.
- CLINTON, B. 2011. Stream water responses to timber harvest: Riparian buffer width effectiveness. *For. Ecol. Manage.* 261:979–988.
- DISSMEYER, G.E. 2000. *Drinking water from forests and grasslands: A synthesis of the scientific literature*. USDA For. Serv., Gen. Tech. Rep. SRS-39, Southern Research Station, Asheville, NC. 246 p.
- DREPS, C., A.L. JAMES, G. SUN, AND J.L. BOOGS. 2014. Water balances of two Piedmont headwater catchments: Implications for regional hydrologic landscape classification. *J. Am. Water Resour. Assoc.* 50(4):1063–1079.
- FRASER, N., R. JACKSON, AND D. RADCLIFFE. 2012. A paired watershed investigation of silvicultural best management practices revisited: B.F. Grant Memorial Forest, Georgia. *For. Sci.* 58(6):652–662.
- GRANT, G., S. LEWIS, F. SWANSON, J. CISSEL, AND J. McDONNELL. 2008. *Effects of forest practices on peak flows and consequent channel response: A state-of-science report for western Oregon and Washington*. USDA For. Serv., Gen. Tech. Rep. PNW-GTR-760, Pacific Northwest Research Station, Portland, OR. 76 p.
- GREENBURG, A.E. 1992. *Standard methods for the examination of water and wastewater*, 18th ed. American Public Health Association, Washington, DC. 1100 p.

- GRIZZEL, J.D., AND N. WOLFF. 1998. Occurrence of windthrow in forest buffer strips and its effect on small streams in Northwest Washington. *Northw. Sci.* 72(3):241–223.
- HEWLETT, J.D., AND A.R. HIBBERT. 1967. Factors affecting the response of small watersheds to precipitation in humid areas. P. 275–290 in *Forest hydrology*, Sopper, W.E., and H.W. Lull (eds.). Pergamon Press, New York.
- HEWLETT, J.D. 1979. Forest water quality: An experiment in harvesting and regenerating Piedmont forests. University of Georgia, Georgia Forest Research Paper, Athens, GA. 22 p.
- HORNBARGER, G.M., K.E. BENCALA, AND D.M. MCKNIGHT. 1994. Hydrological controls on dissolved organic carbon during snowmelt in the Snake River near Montezuma, Colorado. *Biogeochemistry* 2(5):147–165.
- JACKSON, C., D. BATZER, S. CROSS, S. HAGGERTY, AND C. STURM. 2007. Headwater streams and timber harvest: Channel, macroinvertebrate, and amphibian response and recovery. *For. Sci.* 53(2):356–370.
- JACKSON, C.R., G. SUN, D. AMATYA, W.T. SWANK, M. RIEDEL, J. PATRIC, T. WILLIAMS, ET AL. 2004. Fifty years of forest hydrology in the Southeast. P. 33–112 in *A century of forested and wildland watershed lessons*, Ice, G.G., and J.D. Stednick. The Society of American Foresters, Bethesda, MD.
- JOHNSON, E.A., AND J.L. KOVNER. 1954. Increasing water yield by cutting forest vegetation. *Georgia Miner Newsl.* 7:145–148.
- KOCHENDERFER, J.N., P.J. EDWARDS, AND F. WOOD. 1997. Hydrologic impacts of logging an Appalachian watershed using West Virginia's best management practices. *North. J. Appl. For.* 14:207–218.
- LENAT, D.R. 1993. A biotic index for the southeastern United States: Derivation and list of tolerance values, with criteria for assigning water quality ratings. *J. North Am. Benthol. Soc.* 12:279–290.
- LENAT, D.R. 2014. *Biological monitoring of Bolin Creek and Tributaries, Carrboro, North Carolina*. Lenat Consulting Services, Raleigh, NC. 32 p.
- LOCKABY, B.G., F.C. THORNTON, R.H. JONES, AND R.G. CLAWSON. 1994. Ecological response of an oligotrophic floodplain forest to harvesting. *J. Environ. Qual.* 23:901–906.
- MCBROOM, M.W., R.S. BEASLEY, M. CHANG, AND G.G. ICE. 2008. Storm runoff and sediment losses from forest clearcutting and stand re-establishment with best management practices in East Texas, USA. *Hydrolog. Proc.* 22: 1509–1522.
- MOORE, M.K. 1977. *Factors contributing to windthrow in streamside leave strips on Vancouver Island*. Canada Land Management Rep. 3, British Columbia Ministry of Forest, Victoria, BC, Canada. 34 p.
- NORTH CAROLINA DEPARTMENT OF ENVIRONMENT AND NATURAL RESOURCES. 2012. *Standard operating procedures for benthic macroinvertebrates*. Division of Water Quality, Biological Assessment Unit, Raleigh, NC. 49 p.
- PEET, R.K., T.R. WENTWORTH, AND P.S. WHITE. 1998. A flexible, multipurpose method for recording vegetation composition and structure. *Castanea* 63(3):262–274.
- POST, D.A., AND J.A. JONES. 2001. Hydrologic regimes of forested, mountainous, headwater basins in New Hampshire, North Carolina, Oregon, and Puerto Rico. *Adv. Water Resour.* 24:1195–1210.
- RAWER-JOST, C., J. BOHMER, J. BLANK, AND H. RAHMANN. 2004. Macroinvertebrate functional feeding group methods in ecological assessment. *Hydrobiologia* 442/443:225–232.
- RIVENBARK, B.L., AND C.R. JACKSON. 2004. Concentrated flow breakthroughs moving through silvicultural streamside management zones: Southeastern Piedmont, USA. *J. Am. Water Resour. Assoc.* 40(4):1043–1052.
- SAS INSTITUTE, INC. 2011. *JMP*, version 11. SAS Institute, Inc., Cary, NC.
- SINTON, D.S., J.A. JONES, J.L. OHMANN, AND F.J. SWANSON. 2000. Windthrow disturbance, forest composition, and structure in the Bull Run Basin, Oregon. *Ecology* 81(9): 2539–2556.
- STEDNICK, J.D. 1996. Monitoring the effects of timber harvest on annual water yield. *J. Hydrol.* 176:79–95.
- STEIL, J.C., C.R. BLINN, AND R. KOLKA. 2009. Foresters' perceptions of windthrow dynamics in northern Minnesota riparian management zones. *North. J. Appl. For.* 26(2):76–82.
- STEINBLUMS, I.J., H.A. FROELICH, AND J.K. LYONS. 1984. Designing stable buffer strips for stream protection. *J. For.* 82(1):49–52.
- SUN, G., S.G. MCNULTY, D.M. AMATYA, R.W. SKAGGS, L.W. SWIFT JR., J.P. SHEPARD, AND H. RIEKERK. 2002. A comparison of the watershed hydrology of coastal forested wetlands and the mountainous uplands in the southern US. *J. Hydrol.* 263:92–104.
- SUN, G., M. RIEDEL, R. JACKSON, R. KOLKA, D. AMATYA, AND J. SHEPARD. 2004. Influences of management of Southern forests on water quantity and quality. P. 195–234 in *Southern forest science: Past, current, and future*. USDA For. Serv., Gen. Tech. Rep. SRS-GTR-75, Southern Research Station, Asheville, NC.
- SWANK, W.T., AND D.A. CROSSLEY. 1988. *Forest hydrology and ecology at Coweeta*. Ecological Studies 66, Springer-Verlag, New York. 16 p.
- SWANK, W.T., AND J.M. VOSE. 1994. Long-term hydrologic and stream chemistry responses of southern Appalachian catchments following conversion from mixed hardwoods to white pine. P. 164–172 in *Hydrologie kleiner Einzugsgebiete: Gedenkschrift Hans M. Keller*, Landolt, R. (ed). Schweizerische Gesellschaft für Hydrologie und Limnologie, Bern, Switzerland.
- SWANK, W.T., J.M. VOSE, AND K.J. ELLIOTT. 2001. Long-term hydrologic and water quality responses following commercial clear-cutting of mixed hardwoods on a southern Appalachian catchment. *For. Ecol. Manage.* 143:163–178.
- US DEPARTMENT OF AGRICULTURE, SOIL CONSERVATION SERVICE. 1971. *Guide for interpreting engineering uses of soils*. US Government Printing Office, Washington, DC. 86 p.
- US ENVIRONMENTAL PROTECTION AGENCY. 1993. *Paired watershed study design*. EPA No. 841-F-93-009, Office of Water, Washington, DC. 8 p.
- WILLIAMS, T.M., D.D. HOOK, D.J. LIPSCOMB, X. ZENG, AND J.W. ALBISTON. 2000. Effectiveness of best management practices to protect water quality in South Carolina Piedmont. P. 271–277 in *Proc. of the tenth biennial southern silvicultural research conference*, Waldrop, A. (ed.). USDA For. Serv., Gen. Tech. Rep. SRS-30T, Southern Research Station, Asheville, NC.
- WILM, H.G. 1944. Statistical control of hydrologic data from experimental watersheds. *Trans Am. Geophys. Union* Pt. 2:616–622.
- WILM, H.G. 1948. *How long should experimental watershed be calibrated?* USDA For. Serv., Res. Note 2, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 7 p.
- WYNN, T., AND S. MOSTAGHIMI. 2006. The effects of vegetation and soil type on streambank erosion, southwestern Virginia, USA. *J. Am. Water Resour. Assoc.* 42(1):69–82.